



US011285201B2

(12) **United States Patent**
Locht et al.

(10) **Patent No.:** **US 11,285,201 B2**
(45) **Date of Patent:** **Mar. 29, 2022**

(54) **ATTENUATED *BORDETELLA* STRAINS**

(71) Applicants: **Institut Pasteur de Lille**, Lille (FR);
**Institut National de la Santé et de la
Recherche Médicale (INSERM)**, Paris
(FR)

11,110,161 B2 9/2021 Locht et al.
2019/0175719 A1 6/2019 Locht
2019/0290903 A1 9/2019 Zarafshani
2020/0171169 A1 6/2020 Duvall
2020/0206331 A1 7/2020 Kumar
2020/0297833 A1 9/2020 Debrie
2021/0290667 A1 9/2021 Solans

(72) Inventors: **Camille Locht**, Brussels (BE); **Nathalie
Mielcarek**, Blandain (BE);
Anne-Sophie Debrie, La Madeleine
(FR); **Dominique Raze**, Gruson (FR);
Julie Bertout, Pont-à-Marcq (FR)

FOREIGN PATENT DOCUMENTS

EP 1184459 3/2002
WO WO 03/102170 * 12/2003
WO 2008118592 10/2008

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

OTHER PUBLICATIONS

(21) Appl. No.: **16/787,767**

(22) Filed: **Feb. 11, 2020**

(65) **Prior Publication Data**

US 2020/0171139 A1 Jun. 4, 2020

Related U.S. Application Data

(60) Continuation of application No. 16/278,562, filed on
Feb. 18, 2019, now Pat. No. 10,610,580, which is a
division of application No. 15/626,331, filed on Jun.
19, 2017, now Pat. No. 10,258,681, which is a
continuation of application No. 14/860,210, filed on
Sep. 21, 2015, now Pat. No. 9,730,995, which is a
continuation of application No. 14/658,817, filed on
Mar. 16, 2015, now Pat. No. 9,180,178, which is a
division of application No. 12/224,895, filed as
application No. PCT/EP2007/001942 on Mar. 7,
2007, now Pat. No. 9,119,804.

(60) Provisional application No. 60/817,430, filed on Jun.
30, 2006, provisional application No. 60/780,827,
filed on Mar. 10, 2006.

(51) **Int. Cl.**
A61K 39/02 (2006.01)
A61K 39/00 (2006.01)

(52) **U.S. Cl.**
CPC **A61K 39/099** (2013.01); **A61K 2039/522**
(2013.01); **A61K 2039/523** (2013.01); **A61K**
2039/543 (2013.01)

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,883,761 A 11/1989 Keith et al.
9,415,077 B2 8/2016 Alonso et al.
9,528,086 B2 12/2016 Locht et al.
9,655,959 B2 5/2017 Alonso et al.
10,653,765 B2 5/2020 Locht et al.
10,682,377 B2 6/2020 Solans et al.
10,751,072 B2 8/2020 Kendall
10,799,573 B2 10/2020 Brickman et al.
11,065,276 B2 7/2021 Solans et al.

English translation of Debrie et al, 19 pages.*
Mattoo et al, *Frontiers in Bioscience* 6:e168-e186, 2001.*
Gross, Mary K. et al: "Targeted mutations that ablate either the
adenylate cyclase or hemolysin function of the bifunctional cyaA
toxin of *Bordetella pertussis* abolish virulence," *Proc. Natl. Acad.
Sci. USA*, Jun. 1992, vol. 89: 4898-4902.
Lim, Annabelle et al.: "Protective role of adenylate cyclase in the
context of a live pertussis vaccine candidate," *Microbes and Infec-
tion*, 2014, vol. 16:51-60.
Wang, Xianzhe and Jennifer A. Maynard: "The *Bordetella* Adenylate
cyclase repeat-in-toxin (RTX) domain is immunodominant and
elicits neutralizing antibodies," *The Journal of Biological Chemis-
try*, 2015, vol. 290, No. 6:3576-3591.
Storsaeeter, Jann and Joanne Wolter: "Is there a need for a new
generation of vaccines against pertussis?", *Expert Opinion on
Emerging Drugs*, 2006, vol. 11 (2): 195-205.
Scanlon, Karen et al.: "Role of major toxin virulence factors in
pertussis infection and disease pathogenesis," *Adv Exp Med Biol.*,
2019, 1183:: 35-51.
Romero, Rodrigo Villarino et al.: "Filamentous hemagglutinin of
Bordetella pertussis: a key adhesin with immunomodulatory properties?",
Future Microbiol., 2014, vol. 9 (12): 1339-1360.
Poulain-Godefroy, Odile et al.: "Bordetella pertussis filamentous
hemagglutinin delivered by mucosal routes enhances immunoglobulin
levels in serum and mucosal fluids," *FEMS Immunol Med
Microbiol*, 2008, No. 54:129-136.
Melvin, Jeffrey A. et al.: "Bordetella pertussis pathogenesis: current
and future challenges," *Nat Rev Microbiol*, Apr. 2014, vol. 12(4):
274-288.
Li, rui et al/: "Attenuated bordetella pertussis BPZE1 as a live
vehicle for heterologous vaccine antigens delivery through the nasal
route," *Bioengineered Bugs*, 2011, No. 2:6:315-319:449-486.
Kilgore, Paul E. et al: "Pertussis: Microbiology, Disease, Treatment,
and Prevention," *Clinical Microbiology Reviews*, Jul. 2016, vol. 29,
No. 3.

(Continued)

Primary Examiner — Patricia Duffy

(74) *Attorney, Agent, or Firm* — Stanley A. Kim

(57) **ABSTRACT**

A mutated *Bordetella* strain comprising at least a mutated
ptx gene, a deleted or mutated dnt gene and a heterologous
ampG gene is provided. The attenuated mutated *Bordetella*
strain can be used in an immunogenic composition or a
vaccine for the treatment or prevention of a *Bordetella*
infection. Use of the attenuated *Bordetella* strain for the
manufacture of a vaccine or immunogenic composition, as
well as methods for protecting mammals against infection
by *Bordetella* are also provided.

4 Claims, 14 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited

OTHER PUBLICATIONS

Carbonetti, Nicholas H.: "Pertussis toxin and adenylate cyclase toxin: key virulence factors of *Bordetella pertussis* and cell biology tools," *Future Microbiol*, Mar. 2010, vol. 5: 455-469.

Carbonetti, Nicholas H.: "Contribution of pertussis toxin to the pathogenesis of pertussis disease," *FEMS Pathogens and Disease*, Sep. 2015, vol. 73, No. 8:1-8.

* cited by examiner




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10

Islet-activating protein S1 (NP_882282)

MRCTRAIRQTARTGWLTWLAAILAVTAPVTSPAWADDPPATVYRYDSRPPEDVF
QNGFTAWGNNDNVLDHLTGRSCQVGSSNSAFVSTSSSRRYTEVYLEHRMQEAV
EAERAGRGTGHFIGYIYEVRADNNFYGAASSYFEYVDITYGDNAGRILAGALAT
YQSEYLAHRRIPPENIRRVTRVYHNGITGETTTTEYSNARYVVSQQTRANPNPY
TSRRSVASIVGTLVRMAPVIGACMARQAESSEAMAASERAGEAMVLVYYESI
AYSF

Fig. 11

Dermonecrotic toxin (NP_881965)

MDKDESALRQLVDMALVGYDGVVEELLALPSEESGDLAGGRAKREKAEFALFS
EAPNGDEPIGQDARTWFYFPKYRPVAVSNLKKMQVAIRARLEPESLILQWLIA
LDVYLGVLIAALSRTVISDLVFEYVKARYEIIYLLNRVPHPLATAYLKRRRQR
PVDRSGRLGVSFEHPLWFAYDELAGTVDLADIEQALAESIERRMDGEPDDG
SLDTAEHDVWRLCRDGINRGEQAIFQASGPYGVVADAGYMRTVADLAYADALA
DCLHAQLRIRAQGSVDS PGDEMPRKLDWEIAKFHLAATQQARVDLLEAAFAL
DYAALRDVRVYGDYRNALALRFIKREALRLLGARRGNASTMPAVAAGEYDEIV
ASGAANDAAYVSMAAALIAGVLCDESAQRTLPPVVLARFRPLGVLARFRRLAQ
ETAGMLLDGQEPPEPRGFISFTDFRDSDAFASYAEYAAQFNDYIDQYSILEAQR
LARILALGSRMTVDQWCLPLQKVRHYKVLTSQPGLIARGIENHNRGIEYCLGR
PPLTDLPLGLFTMFQLHDSSWLLVSNINGELWSDVLANAEMQNPTLAALAEPO
GRFRTGRRRTGGWFLGGPATEGPSLRDNYLLKLRQSNPGLDVKKCWYFGYRQEI
RLPAGALGVPLFAVSVALRHSLDDLAHAHAKSALYKPSWQKFAFWIVPFYREI
FFSTQDRSYRVDVGSIVFDSISLLASVFSIGGKLGSTRTQYGNLRFVVRQR
IAGLSGQRLWRSVLKELPALIGASGLRLSRSLLDVLYEIFEPVPIRRLVAGFV
SATTVGGRNQAFRLRQAFSAASSAGRTGGQLASEWRMAGVDATGLVESTSGGR
FEGIYTRGLGPLSECTEHFIVESGNAYRVIWDAYTHGWRVNVNRLPPRLTYTV
PVRLNGQGHWETHLDVPGRGGAPEIFGRIRTRNLVALAAEQAAPMRLLNQAR
RVALRHIDTCRSRLALPRAESDMDAAIRIFFGEPDAGLRQRIGRRLQEVRAI
GDLSPVNDVLYRAGYDLDDVATLFNAVDRNTSLGRQARMELYLDAIVDLHARL
GYENARFVDLMAFHLLSLGHAATASEVVEAVSPRLLGNVFDISNVAQLERGIG
NPASTGLFVMLGAYSESSPAIFQS FVNDIFPAWRQASGGGPLVWNFGPAAISP
TRLDYANTDIGLLNHGDISPLRARPPPLGGRRDIDLPPGLDISFVRYDRPVRMS
APRALDASVFRPVDGPHVGYIQSWTGAEIEYAYGAPAAAREVMLTDNVRIISI
ENGDEGAIGVRVRLDTPVATPLILTGGSLSGCTTMVGVKEGYLAFYHTGKST
ELGDWATAREGVQALYQAHLAMGYAPISIPAPMRNDDLVSIAATYDRAVIAYL
GKDVPGGGSTRITRHDEGAGSVVSFDYNAAVQASAVPRLGQVYVLI SNDGQGA
RAVLLAEDLAWAGSGSALDVLNERLVTLFPAPV

Fig. 12

AmpG protein (NP_878961.1)

MAPLLVLGFASGLPLALSSGTLQAWATVENVSLQSIGFLTLAGTAYTLKFLWA
PLIDRYVPPFLGRRRGWMLLTQVLLAAAIMVMGMLSPGSALLPLALVAVLVA
LSASQDIAFDAYSTDVLRQEERGAGAAMRVMGYRLAMIVSGGLALIVADRWL
WGNTYVLMGGLMLACALGTLWAPEPERPANPPRDLGAAVVEPFREFFSRRGAI
DMLLLIVLYKLGDAFAGALSTTFLLRGAGFSATEVGTVNKVLGLAATIVGALA
GGSIMTRWGLYRSLMAFGLLQAVSNLGYWLI AVSPKNLYLMGLAVGVENLCGG
LGTASFVALLMAMCRQQFSATQFALLSALAAVGRTYLAGPLTPVLVEWLDWPG
FFIVTVLIALPGLWLLRLRRNVIDELDAQAR

Fig. 13

AmpG protein (NP_752478.1)

MSSQYLRIFQQPRSAILLILGFASGLPLALTSQTLQAWMTVENIDLKTIGFFS
LVGQAYVFKFLWSPLMDRYTPPFFGRRRGWLLATQILLLVIAAAMGFLEPGTQ
LRWMAALAVVIAFCSASQDIVFDAWKTDVLPAEERGAGAAISVLGYRLGMLVS
GGLALWLADKWLGWQGMWYLMALLIPCI IATLLAPEPTDTIPVPKTLEQAVV
APLRDFFGRNNAWLILLIVLYKLGDAFAMSLTTTFLIRGVGFDAGEVGVVVK
TLGLLATIVGALYGGILMQRSLFRALLIFGILQGASNAGYWLLSITDKHLYS
MGAAVFFENLCGGMGTSAFVALLM TLCNKSFSATQFALLSALS AVGRVYVGPV
AGWFVEAHGWSTFYLF SVAAAVPGLILLLVCRQTLEYTRVNDNFISRTEYPAG
YAFAMWTLAAGISLLAVWLLLLTMDALDLTHFSFLPALLEVGVLVALS GVVLG
GLLDYLALRKTHLM

Fig. 14

ATTENUATED *BORDETELLA* STRAINS

CROSS-REFERENCE TO RELATED APPLICATIONS

The present application is a continuation application of U.S. patent application Ser. No. 16/278,562 filed on Feb. 18, 2019, which is a division of U.S. patent application Ser. No. 15/626,331 filed on Jun. 19, 2017 (now U.S. Pat. No. 10,258,681), which is a continuation application of U.S. patent application Ser. No. 14/860,210 filed on Sep. 21, 2015 (now U.S. Pat. No. 9,730,995), which is a continuation application of U.S. patent application Ser. No. 14/658,817 filed on Mar. 16, 2015 (now U.S. Pat. No. 9,180,178), which is a divisional application of U.S. nonprovisional patent application Ser. No. 12/224,895 filed on Nov. 19, 2008 (now U.S. Pat. No. 9,119,804) as a national stage entry application under 35 U.S.C. 371 of international patent application number PCT/EP2007/01942, filed on Mar. 7, 2007, which designated the U.S. and claims the priority of U.S. provisional patent application Ser. No. 60,817,430 filed on Jun. 30, 2006 and U.S. provisional patent application Ser. No. 60/780,827 filed on Mar. 10, 2006, which are incorporated herein by reference in their entirety.

FIELD OF THE INVENTION

The present invention relates to a mutated *Bordetella* strain comprising at least a mutated *ptx* gene, a deleted or mutated *dnt* gene and a heterologous *ampG* gene. The attenuated mutated *Bordetella* strain can be used in an immunogenic composition or a vaccine for the treatment or prevention of a *Bordetella* infection. Use of the attenuated *Bordetella* strain for the manufacture of a vaccine or immunogenic compositions, as well as methods for protecting mammals against infection by *Bordetella* also form a part of the invention.

BACKGROUND OF THE INVENTION AND RELATED PRIOR ART

Pertussis is still among the principal causes of death world-wide, and its incidence is increasing even in countries with high vaccine coverage. Although all age groups are susceptible, it is most severe in infants too young to be protected by currently available vaccines.

Whooping cough or pertussis is a severe childhood disease responsible for high mortality rates before the introduction of effective vaccines in the second half of the 20th century. The success of these vaccines has led to the opinion that the disease is essentially under control, although world-wide 200,000 to 400,000 pertussis-linked deaths are still recorded annually, and the disease still ranks sixth among the causes of mortality due to infectious agents [1]. Although mostly prevalent in developing countries, the disease is also re-emerging in the developed world [2, 3], including the U.S.A., where the incidence has increased five-fold over the last twenty years [4]. Unexpectedly, the epidemiology of pertussis has changed in countries with high vaccine coverage, where cases of adolescent and adult pertussis are increasingly frequent [5]. This is probably due to progressive waning of vaccine-mediated immunity during adolescence. Often atypical and therefore difficult to diagnose, pertussis is generally not life-threatening in adults and in many cases remains unnoticed. However, infected adults constitute an important reservoir for transmission of the disease to very young children, too young to be fully

vaccinated, and therefore at risk to develop severe disease associated with high mortality rates.

Pertussis vaccination usually begins at two months of age, and full protection requires at least three immunizations at one- to two-month intervals. Therefore, infants are not fully protected before the age of 6 months using the currently available vaccines. To reduce the incidence of pertussis in the very young and most vulnerable age groups, early immunization, possibly at birth, would thus be highly desirable. However, numerous studies in humans and in animal models have suggested that the neonatal immune system is too immature to effectively induce vaccine-mediated protective immunity [6, 7]. Especially the IFN- γ production, indicative of a Th1 response that is essential to the development of protective immunity to pertussis [8], appears to be significantly reduced in human newborns, compared to older children or adults [9]. This is also reflected by the fact that significant amounts of antigen-specific IFN- γ are only produced after several months (≥ 6 months) in children vaccinated with pertussis vaccines, especially with acellular vaccines (aPV) [10].

Natural infection with *Bordetella pertussis* has long been considered to induce strong and long-lasting immunity, that wanes much later than vaccine-induced immunity [5, 11]. Furthermore, infection with *B. pertussis* induces measurable antigen-specific Th1 type immune responses even in very young children (as young as one month of age) [12]. These observations suggest that live vaccines applicable by the nasal route in order to mimic as closely as possible natural infection, may be attractive alternatives over the currently available vaccines.

There are many vaccinating compositions to treat *Bordetella* infections known in the art. However, these immunogenic compositions are not used to treat newborn children or in cases where an epidemic and rapid protective immunity is required.

Thus, French Patent FR 0206666 discloses live *Bordetella* strains that have been rendered deficient in at least two toxins chosen from PTX, DNT, AC and TCT. This patent discloses the over expression of an endogenous *ampG* gene by the addition of a strong promoter, and the addition of 11 terminal amino acids of the *ampG* gene from *E. coli*.

Mielcarek et al, Vaccine (2006; 24(52): 52154-52-55) disclose a strain of *Bordetella pertussis* attenuated of PTX, DTN- and TCr for use in the immunization of mice. This reference discloses that to reduce the production of tracheal cytotoxin, the *ampG* gene should be overexpressed. However, upon further evaluation, the authors realized that by over-expressing the *ampG* gene, there is an increase in tracheal cytotoxin and not a decrease as was originally thought.

Mielcarek et al in Advance Drug Delivery Review 51 (2001) pgs. 55-69 disclose that live vaccines can induce systemic and mucosal responses when administered by the oral or nasal route.

Roduit et al in Infection and Immunity (2002 July; 70(7): 3521-8) describe vaccinating neonatals and infants with mutated *Bordetella* strains with a DTP composition.

Mattoo et al, in Frontiers of Bioscience 6, e168-e186 (2001), suggest replacing the endogenous *ampG* gene in *Bordetella* with the *E. coli ampG* gene, which resulted in a decrease in the amount of TCT produced.

Thus, the prior art although disclosing various types of vaccinating compositions fails to address the problem of providing a vaccine or immunogenic composition that can provide protection to a newborn prior to six months. Furthermore, the prior art fails to disclose an immunogenic or

a vaccine that provides rapid protective immunity against a *Bordetella* infection. The prior art also fails to disclose an immunogenic composition or vaccine that provides a rapid protective immunity against a *Bordetella* infection, said protective immunity increasing over at least the next two months following vaccination.

Therefore, it is an object of the present invention to overcome the deficiencies in the prior art.

It is another object of the present invention to produce a live attenuated vaccine candidate or immunogenic composition through genetic attenuation of a *Bordetella* strain such as *B. pertussis* or *B. parapertussis* to diminish pathogenicity, while maintaining the ability to colonize and induce protective immunity.

It is another object of the present invention to produce a vaccine or immunogenic composition that induces protection in newborns after a single intranasal administration that is superior to the protection provided by the current aPV.

It is yet another object of the present invention to provide protection against infection with *Bordetella parapertussis*, as well as *Bordetella pertussis* which was not seen after vaccination with aPV.

Another object of the present invention is to induce strong protective immunity in newborns against *Bordetella* infection.

Yet another object of the present invention is to provide a vaccine or immunogenic composition that induces mucosal and systemic immunity.

It is another object of the present invention to produce a live attenuated *Bordetella pertussis* strain to be given as a single-dose nasal vaccine in early life, called BPZE1.

It is yet another object of the present invention to provide a vaccine that can not only be used to vaccinate newborns, but can be used in all mammals of any age in the case of an epidemic of whooping cough.

Another object of the present invention is to provide a vaccine against *Bordetella* infection that induces a rapid protective immunity and/or a protective immunity that increases over at least the next two months after the vaccination.

Yet another object of the present invention is to provide prevention or treatment against *Bordetella* infection that is relatively low in production costs.

These and other objects are achieved by the present invention as evidenced by the summary of the invention, description of the preferred embodiments and the claims.

SUMMARY OF THE INVENTION

The present invention provides a mutated *Bordetella* strain comprising at least a mutated pertussis toxin (ptx) gene, a deleted or mutated dermonecrotic toxin (dnt) gene, and a heterologous ampG gene.

In another aspect the present invention relates to an immunogenic composition comprising a mutated *Bordetella* strain comprising at least a mutated pertussis toxin (ptx) gene, a deleted or mutated pertussis dermonecrotic toxin (dnt) gene, and a heterologous ampG gene.

In yet another aspect the present invention provides a vaccine comprising the attenuated *Bordetella* strain comprising at least a mutated pertussis toxin (ptx) gene, a deleted or mutated pertussis dermonecrotic toxin (dnt) gene, and a heterologous ampG gene.

In still another aspect, the present invention provides the use of an attenuated *Bordetella* strain comprising at least a mutated ptx gene, a deleted or mutated dnt gene, and a

heterologous ampG gene for the manufacture of a vaccine for the prevention of a *Bordetella* infection.

In yet another aspect, the present invention provides the use of an attenuated *Bordetella* strain comprising at least a mutated ptx gene, a deleted or mutated dnt gene, and a heterologous ampG gene for the manufacture of a vaccine for the induction of an immune response directed preferentially toward the Th1 pathway against said attenuated *Bordetella*.

Also provided is a method of protecting a mammal against disease caused by infection by *Bordetella pertussis* and *Bordetella parapertussis* comprising administering to said mammal in need of such treatment a mutated *Bordetella* strain comprising at least a mutated ptx gene, a deleted or mutated dnt gene, and a heterologous ampG gene.

A method of providing a rapid protective immunity against a *Bordetella* infection comprising administering to said mammal in need of such treatment a mutated *Bordetella* strain comprising at least a mutated ptx gene, a deleted or mutated dnt gene, and a heterologous ampG gene is also part of the present invention.

A method of providing a rapid protective immunity against a *Bordetella* infection comprising administering to a mammal in need of such treatment a mutated *Bordetella* strain comprising at least a mutated ptx gene, a deleted or mutated dnt gene, and a heterologous ampG gene or a vaccine comprising said mutated *Bordetella* strain, wherein said method provides further an increase in said protective immunity over at least two months after vaccination is still another aspect of the present invention.

Use of the mutated *Bordetella* strain comprising at least a mutated ptx gene, a deleted or mutated dnt gene and a heterologous ampG gene for the preparation of a multivalent vaccine (i.e., a vaccine for preventing or treating infections caused by different pathogens) to treat respiratory diseases is yet another aspect of the present invention.

Use of an attenuated *Bordetella* strain of the invention, by administration to mammals in need of a rapid protective immunity against a *Bordetella* infection, wherein said protective immunity increases over at least two months after administration, is also part of the present invention.

A method to provide a mucosal response and a systemic response to treat or protect against *Bordetella* infections in mammals is still another aspect of the present invention.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a bar graph illustrating the TCT present in culture supernatants of BPSM and BPZE1 expressed as means of $\text{nM}/\text{OD}_{540 \text{ nm}}$ (\pm standard error) of 3 separate cultures for each strain.

FIG. 2 is an immunoblot analysis of PTX production in the culture supernatants of BPSM (lane 1) and BPZE1 (lane 2). The sizes of the Mr markers are expressed in kDa and given in the left margin.

FIG. 3 is a Southern-blot analysis of the dnt locus in BPSM (lane 1) and BPZE1 (lane 2). The lengths of the size markers are indicated in base pairs (bp) are shown in the left margin.

FIG. 4 is a graph illustrating the growth rates of BPSM (black line) and BPZE1 (dotted line) in liquid culture.

FIG. 5 are electron micrographs representative of BPSM (left) and BPZE1 (right) grown in liquid medium for 24 h.

FIG. 6 is a graph illustrating the in vitro adherence of BPSM (black columns) and BPZE1 (white columns) to human pulmonary epithelial A549 cells (left) and murine macrophage-like J774 cells (right). The results are expressed

as means of percentages of binding bacterial relative to the bacteria present in the inoculum from three different experiments.

FIG. 7 is a graph illustrating lung colonization by BPSM (black lines) and BPZE1 (dotted lines) of adult mice infected intranasally with 10⁶ CFU of BPZE1 or BPSM. The results are expressed as mean (\pm standard error) CFUs from three to four mice per group and are representative of two separate experiments. P=0.004.

FIG. 8 are photographs of a histological analysis of lungs from BPZE1 (upper panel) or BPSM-infected (middle panel) adult mice compared to controls given PBS (lower panel). One week after infection, the lungs were aseptically removed and fixed in formaldehyde. Sections were stained with hematoxylin and eosin and examined by light microscopy.

FIG. 9 are graphs illustrating the protection against *B. pertussis* in (a) adult and (b) infant mice or *B. parapertussis* in infant mice (d). Mice immunized with BPZE1, aPV or PBS (naive) were challenged with BPSM (a and b) or *B. parapertussis* (d), and lung CFU counts were determined 3 h (white bars) or 7 days (black bars) later. Results are expressed as mean (\pm standard error) CFUs from 3-4 mice per group and are representative of two separate experiments. (b, *, P=0.009; d, *, P=0.007) (c) CFU counts 3 h after BPSM challenge in adult mice vaccinated with BPZE1 or aPV, compared to controls. Results obtained from 3 separate experiments are expressed as percentages of CFUs of each mouse relative of the average of CFUs in non-immunized group from the same experiment.

FIG. 10 are bar graphs illustrating the immune responses induced by BPZE1 or aPV immunization. (a) Anti-FHA IgG(H+I) titers and (b) IgG1/IgG2a ratios before (white bars) or 1 week after BPSM challenge (black bars) in BPZE1 or aPV immunized mice, compared to controls. (c) IFN- γ to IL-5 ratios produced by FHA-, PTX- or ConA-stimulated splenocytes from mice vaccinated 2 months before with BPZE1 (black bars) or aPV (white bars), compared to controls (gray bars). Antibodies and cytokines were measured in individual mice, and the results are expressed as mean values (\pm standard error) for 4 mice per group tested in triplicate.

FIG. 11 is the amino acid sequence of pertussis toxin (SEQ 10 NO:1) (islet-activating protein 51). The first 34 amino acids are the signal sequence, while amino acids 35 to 269 are the mature chain.

FIG. 12 is the amino acid sequence of dermonecrotic toxin (SEQ ID NO:2).

FIG. 13 is the amino acid sequence of AmpG from *Bordetella pertussis* (SEQ ID NO:3).

FIG. 14 is the amino acid sequence of AmpG from *Escherichia coli* (SEQ ID NO:4).

DESCRIPTION OF THE PREFERRED EMBODIMENTS OF THE PRESENT INVENTION

As used herein, the abbreviation "PTX" refers to pertussis toxin, which synthesizes and secretes an ADP-ribosylating toxin. PTX is composed of six polypeptides S1 to S5, the enzymatically active moiety is called S1. PTX has a 34 amino acid signal sequence, while the mature chain consists of amino acids 35 to 269. PTX is the major virulence factor expressed by *B. pertussis*. The A moiety of these toxins exhibit ADP-ribosyltransferase activity and the B portion mediates binding of the toxin to host cell receptors and the translocation of A to its site of action (57).

As used herein the abbreviation "DNT" refers to pertussis dermonecrotic toxin, which is a heat labile toxin that induces localized lesions in mice and other laboratory animals when it is injected intradermally. It is lethal to mice when it is injected in low doses intravenously (58 to 61). DNT is considered to be a virulence factor for the production of turbinate atrophy in porcine atrophic rhinitis (62, 63).

As used herein the abbreviation "TCT" refers to tracheal cytotoxin, which is a virulence factor synthesized by *Bordetella*. TCT is a peptidoglycan fragment and has the ability to induce interleukin-1 production and nitric oxide synthase. It has the ability to cause stasis of cilia and has lethal effects on respiratory epithelial cells.

The term "mammal" encompasses any of various warm-blooded vertebrate animals of the class Mammalia, including humans, characterized by a covering of hair on the skin and, in the female, milk-producing mammary glands for nourishing the young.

The term "attenuated" means a weakened, less virulent *Bordetella* strain that is capable of stimulating an immune response and creating protective immunity, but does not cause any illness.

The terminology "rapid protective immunity" means that immunity against *Bordetella* is conferred in a short time after administration of the mutated *Bordetella* strain of the present invention. By "short time" means vaccinated and challenged one week later. More specifically, there is a quick expansion of existing pathogen-specific peripheral lymphocytes, CD8+ cytotoxic effectors (CTLs) and CD4+ helper cells. The CD4+ helper cells induce B cell maturation and antibody production. Thus, lymphocytes with the memory pool are poised to rapidly proliferate at the time of subsequent infection.

The term "*Bordetella* strain" encompasses strains from *Bordetella pertussis*, *Bordetella parapertussis* and *Bordetella bronchiseptica*.

The expression "*Bordetella* infection" means an infection caused by at least one of the three following strains: *Bordetella pertussis*, *Bordetella parapertussis* and *Bordetella bronchiseptica*.

By "child" is meant a person or a mammal between 6 months and 12 years of age.

By the term "newborn" is meant a person or a mammal that is between 1 day old and 24 weeks of age.

The term "treatment" as used herein is not restricted to curing a disease and removing its causes but particularly covered means to cure, alleviate. Remove or lessen the symptoms associated with the disease of interest, or prevent or reduce the possibility of contracting any disorder or malfunction of the host body.

The terms "protection" and "prevention" are used herein interchangeably and mean that an infection by *Bordetella* is impeded.

"Prophylaxis vaccine" means that this vaccine prevents *Bordetella* infection upon future exposure.

By "preferentially towards the Th1 pathway" is meant that the Th1 pathway is favored over the Th2 pathway.

The term "immunogenic composition" means that the composition can induce an immune response and is therefore antigenic. By "immune response" means any reaction by the immune system. These reactions include the alteration in the activity of an organism immune system in response to an antigen and may involve, for example, antibody production, induction of cell-mediated immunity, complement activation or development of immunological tolerance

More specifically, the present invention provides at least a triple mutated *Bordetella* strain that can be used as an immunogenic composition or a vaccine. It will be appreciated that the at least triple mutated *Bordetella* strain contains a mutated ptx gene, a deleted or mutated dnt gene and a heterologous ampG gene. The heterologous ampG gene product reduces in large quantities the amount of tracheal cytotoxin that is produced.

The present invention is not limited to only the triple mutants described above. Other additional mutations can be undertaken such as adenylate cyclase (AC) deficient mutants (64), lipopolysaccharide (LPS) deficient mutants (65), filamentous hemagglutinin (FHA) (66) and any of the bvg-regulated components (67).

The starting strain which is mutated can be any *Bordetella* strain including *Bordetella pertussis*, *Bordetella parapertussis* and *Bordetella bronchiseptica*. In one aspect the starting strain used to obtain the mutated *Bordetella* strain is *B. pertussis*.

The construction of the mutated *Bordetella* strain starts with replacing the *Bordetella* ampG gene in the strain with a heterologous ampG gene. Any heterologous ampG gene can be used in the present invention. These include all those gram-negative bacteria that release very small amounts of peptidoglycan fragments into the medium per generation. Examples of gram-negative bacteria include, but are not limited to *Escherichia coli*, *Salmonella*, *Enterobacteriaceae*, *Pseudomonas*, *Moraxella*, *Helicobacter*, *Stenotrophomonas*, *Legionella* and the like.

By replacing the *Bordetella* ampG gene with a heterologous ampG gene, the amount of tracheal cytotoxin (TCT) produced in the resulting strain expresses less than 1% residual TCT activity. In another embodiment, the amount of TCT toxin expressed by the resulting strain is between 0.6% to 1% residual TCT activity or 0.4% to 3% residual TCT activity or 0.3% to 5% residual TCT activity.

PTX is a major virulence factor responsible for the systemic effects of *B. pertussis* infections, as well as one of the major protective antigens. Due to its properties, the natural ptx gene is replaced by a mutated version so that the enzymatically active moiety S1 codes for an enzymatically inactive toxin, but the immunogenic properties of the pertussis toxin are not affected. This can be accomplished by replacing the arginine (Arg) at position 9 of the sequence with a lysine (Lys). Furthermore, a glutamic acid (Glu) at position 129 is replaced with a glycine (Gly).

Other mutations can also be made such as those described in U.S. Pat. No. 6,713,072, incorporated herein by reference, as well as any known or other mutations able to reduce the toxin activity to undetectable levels. Allelic exchange is first used to delete the ptx operon and then to insert the mutated version.

Finally, the dnt gene is then removed from the *Bordetella* strain by using allelic exchange. Besides the total removal, the enzymatic activity can also be inhibited by a point mutation. Since DNT is constituted by a receptor-binding domain in the N-terminal region and a catalytic domain in the C-terminal part, a point mutation in the dnt gene to replace Cys-1305 to Ala-1305 inhibits the enzyme activity of DNT (68). DNT has been identified as an important toxin in *Bordetella bronchiseptica* and displays lethal activity upon injection of minute quantities (26).

Besides allelic exchange to insert the mutated ptx gene and the inhibited or deleted dnt gene, the open reading frame of a gene can be interrupted by insertion of a genetic sequence or plasmid. This method is also contemplated in the present invention.

The triple mutated strain of the present invention is called a BPZE1 strain and has been deposited with the Collection Nationale de Cultures de Microorganismes (CNCM) in Paris, France on Mar. 9, 2006 under the number CNCM 1-3585. The mutations introduced into BPZE1 result in drastic attenuation, but allow the bacteria to colonize and persist. Thus, in another embodiment the present invention provides BPZE1, which can induce mucosal immunity and systemic immunity when administered. In another aspect the BPZE1 is administered intranasally.

The mutated *Bordetella* strains of the present invention can be used in immunogenic compositions. Such immunogenic compositions are useful to raise an immune response, either an antibody response and or preferably a T cell response in mammals. Advantageously, the T cell response is such that it protects a mammal against *Bordetella* infection or against its consequences.

The mutated *Bordetella* strains of the present invention can be used as live strains or chemically or heat-killed strains in the vaccines or immunogenic compositions. In one aspect, the live strains are used for nasal administration, while the chemically or heat killed strains can be used for systemic or mucosal administration.

The immunogenic composition may further comprise a pharmaceutically suitable excipient or carrier and/or vehicle, when used for systemic or local administration. The pharmaceutically acceptable vehicles include, but are not limited to, phosphate buffered saline solutions, distilled water, emulsions such as an oil/water emulsions, various types of wetting agents sterile solutions and the like.

The immunogenic composition of the invention can also comprise adjuvants, i.e., any substance or compound capable of promoting or increasing a T-cell mediated response, and particularly a CD4⁺-mediated or CD8⁺-mediated immune response against the active principle of the invention. Adjuvants such as muramyl peptides such as MDP, IL-12, aluminium phosphate, aluminium hydroxide, Alum and/or Montanide® can be used in the immunogenic compositions of the present invention.

It would be appreciated by the one skilled in the art that adjuvants and emulsions in the immunogenic compositions are used when chemically or heat treated mutated *Bordetella* strains are used in the vaccines or immunogenic compositions.

The immunogenic compositions of the invention further comprise at least one molecule having a prophylactic effect against a *Bordetella* infection or the detrimental effects of *Bordetella* infection, such as a nucleic acid, a protein, a polypeptide, a vector or a drug.

The immunogenic composition of the invention is used to elicit a T-cell immune response in a host in which the composition is administered. Ail immunogenic compositions described above can be injected in a host via different routes: subcutaneous (s.c.), intradermal (i.d.), intramuscular (i.m.) or intravenous (i.v.) injection, oral administration and intranasal administration or inhalation.

When formulated for subcutaneous injection, the immunogenic composition or vaccine of the invention preferably comprises between 10 and 100 µg of the *Bordetella* strain per injection dose, more preferably from 20 to 60 µg/dose, especially around 50 µg/dose, in a sole injection.

When formulated for intranasal administration, the *Bordetella* strain is administered at a dose of approximately 1×10^3 to 1×10^6 bacteria, depending on the weight and age of the mammal receiving it. In another aspect a dose of 1×10^4 to 5×10^6 can be used.

The mutated *Bordetella* strains of the present invention can be used as an attenuated vaccine to protect against future *Bordetella* infection. In this regard, an advantage of the present invention is that a single dose can be administered to mammals and the protection can last at least for a duration of longer than two months, particularly longer than six months. The vaccine of the present invention can be administered to newborns and protects against infection of whooping cough. This is especially crucial since the fatality rate from *Bordetella pertussis* infections is about 1.3% for infants younger than 1 month.

Moreover, the vaccines of the present invention can be used in adult mammals when there is an epidemic or in older adults over the age of 60, since their risk of complications may be higher than that of older children or healthy adults.

The vaccines can be formulated with the physiological excipients set forth above in the same manner as in the immunogenic compositions. For instance, the pharmaceutically acceptable vehicles include, but are not limited to, phosphate buffered saline solutions, distilled water, emulsions such as an oil/water emulsions, various types of wetting agents sterile solutions and the like. Adjuvants such as muramyl peptides such as MDP, IL-12, aluminium phosphate, aluminium hydroxide, Alum and/or Montanide® can be used in the vaccines.

The vaccines of the present invention are able to induce high titers of serum IgG against FHA. The analysis of the antigen-specific cytokine patterns revealed that administration with the mutated attenuated *Bordetella* strains of the present invention favored a strong TH1 response.

The vaccines of the present invention provide high level of protection against a *Bordetella* infection i.e., a level of protection higher than 90%, particularly higher than 95%, more particularly higher than 99% (calculated 7 days after infection as detailed on example 9). The level of protection of the vaccine comprising the BPZE1 strain reaches more than 99.999% compared to non-vaccinated (naïve) mice, at least two months after vaccination.

The vaccines can be administered subcutaneous (s.c.), intradermal (i.d.), intramuscular (i.m.) or intravenous (i.v.) injection, oral administration and intranasal administration or inhalation. The administration of the vaccine is usually in a single dose. Alternatively, the administration of the vaccine of the invention is made a first time (initial vaccination), followed by at least one recall (subsequent administration), with the same strain, composition or vaccine, or with acellular vaccines, or a combination of both.

In one aspect, intranasal administration or inhalation of the vaccines is accomplished, which type of administration is low in costs and enables the colonization by the attenuated strains of the invention of the respiratory tract: the upper respiratory tract (nose and nasal passages, paranasal sinuses, and throat or pharynx) and/or the respiratory airways (voice box or larynx, trachea, bronchi, and bronchioles) and/or the lungs (respiratory bronchioles, alveolar ducts, alveolar sacs, and alveoli).

Intranasal administration is accomplished with an immunogenic composition or a vaccine under the form of liquid solution, suspension, emulsion, liposome, a cream, a gel or similar such multiphasic composition. Solutions and suspensions are administered as drops. Solutions can also be administered as a fine mist from a nasal spray bottle or from a nasal inhaler. Gels are dispensed in small syringes containing the required dosage for one application.

Inhalation is accomplished with an immunogenic composition or a vaccine under the form of solutions, suspensions, and powders; these formulations are administered via

an aerosol or a dry powder inhaler. Compounded powders are administered with insufflators or puffers.

Use of the mutated *Bordetella* strains comprising at least a mutated ptx gene, a deleted or mutated din gene and a heterologous ampG gene for the preparation of a multivalent vaccine to treat respiratory diseases is yet another aspect of the present invention. In this regard, the attenuated mutated *Bordetella* strain described above, can be used as a heterologous expression platform to carry heterologous antigens to the respiratory mucosa. Thus, such respiratory pathogens such as *Neisseria*, *Pneumophila*, *yersinia*, *pseudomonas*, *mycobacteria*, *influenza* and the like can prevent infection using the BPZE1 as a carrier.

Use of the live attenuated mutated *Bordetella* strains described herein for the manufacture of a vaccine for the treatment or prevention of *Bordetella* infection is also encompassed by the present invention. In this regard, the vaccine can be used for the simultaneous treatment or prevention of an infection by *B. pertussis* and *B. parapertussis*.

Use of the vaccine to provide rapid protective immunity in case of a pertussis epidemic is also encompassed by the present invention.

Use of the vaccine to provide a rapid protective immunity, increasing over the at least next two months following vaccination is also encompassed by the present invention.

The vaccine or immunogenic composition is also provided in a kit. The kit comprises the vaccine or immunogenic composition and an information leaflet providing instructions for immunization.

The present invention also relates to a method for inducing T-cell mediated immune response and particularly a CD4⁺-mediated immune response or a CD8⁺-mediated immune response, comprising administering the live attenuated *Bordetella* strains of the invention in a non-human mammal or a human mammal.

A method of protecting a mammal against disease caused by infection by *Bordetella* comprising administering to said mammal in need of such treatment a mutated *Bordetella* strain comprising at least a mutated ptx gene, a deleted or mutated dnt gene, and a heterologous ampG gene is another embodiment of the present invention. This method encompasses treating or preventing infections against *Bordetella pertussis* and/or *Bordetella parapertussis*. In one aspect the BPZE1 strain is used in this method.

Also a method of providing a rapid protective immunity against a *Bordetella* infection comprising administering to said mammal in need of such treatment a mutated *Bordetella* strain comprising at least a mutated ptx gene, a deleted or mutated dnt gene, and a heterologous ampG gene is encompassed by the present invention. In one aspect the BPZE1 strain is used in this method.

Moreover, the mutated live attenuated *Bordetella* strains of the present invention induce mucosal immunity, as well as systemic immunity. Thus, in another aspect the invention also relates to a method of inducing mucosal and systemic immunity by administering to a mammal in need of such treatment the mutated live attenuated *Bordetella* strains of the present invention. In one aspect the BPZE1 strain is used in this method.

Besides its role in the prevention and/or treatment of *Bordetella* infection, the mutated strain of the invention may be used as vector, to bear at least one further heterologous nucleic acid sequence encoding a RNA (such as antisense RNA) or a protein of interest. This means that the mutated strain bears at least one further heterologous nucleic acid sequence in addition to the heterologous ampG gene. In one

11

aspect, the protein encoded by this at least one further heterologous nucleic acid sequence is a protein for which the expression is desired in the respiratory tract. In another aspect, the protein of interest is an antigen, such as a viral, a bacterial or a tumoral antigen, against which an immune response is desired. Therefore, the mutated *Bordetella* strain bearing at least one further heterologous nucleic acid sequence may also be used as a vaccine. The definitions given above for administration of the vaccine or immunogenic composition also apply to a vaccine comprising mutated *Bordetella* strain bearing at least one further heterologous nucleic acid sequence. Examples of heterologous proteins are antigens of pathogens causing infections of or diseases associated with the respiratory track: poliomyelitis, influenza (influenzavirus from Orthomyxoviridae family) or antigens from pneumococcus (such as *Streptococcus pneumoniae*).

A number of embodiments of the invention have been described. Nevertheless, it will be understood that various modifications may be made without departing from the spirit and scope of the invention.

EXAMPLES

Materials and Methods

Example 1—*Bordetella* Strains and Growth Conditions

The *B. pertussis* strains used in this study were all derived from *B. pertussis* BPSM [13], and *B. parapertussis* is a streptomycin-resistant derivative of strain 12822 (kindly provided by Dr. N. Guiso, Institut Pasteur Paris, France). All *Bordetella* strains were grown on Bordet-Gengou (BG) agar (Difco, Detroit, Mich.) supplemented with 1% glycerol, 20% defibrinated sheep blood, and 100 streptomycin. For cell adherence assays, exponentially growing *B. pertussis* was inoculated at an optical density of 0.15 at 600 nm in 2.5 ml modified Stainer-Scholte medium [14] containing 1 g/l heptakis(2,6-di-*o*-methyl) β -cyclodextrin (Sigma) and supplemented with 65 μ Ci/ml [³⁵S]methionine plus L-[³⁵S]cysteine (NEN, Boston, Mass.) and grown for 24 h at 37° C. The bacteria were then harvested by centrifugation, washed three times in phosphate-buffered saline (PBS) and resuspended in RPMI 1640 (Gibco, Grand Island, N.Y.) at the desired density.

Example 2—Construction of *B. pertussis* BPZE1

To construct *B. pertussis* BPZE1, the *B. pertussis* ampG gene was replaced by *Escherichia coli* ampG using allelic exchange. A PCR fragment named met and located at position 49,149 to 49,990 of the *B. pertussis* genome, upstream of the *B. pertussis* ampG gene, was amplified using oligonucleotides A: 5'-TATAAATCGATATTCCTGCTGGTTTCGTTCTC-3' (SEQ ID No:5) and B: 5'-TATAGCTAGCAAGTTGGGAAACGACACCAC-3' (SEQ ID No 6), and *B. pertussis* BPSM [13] genomic DNA as a template. This 634 bp fragment was inserted into Topo PCR II (Invitrogen Life Technology, Groningen, The Netherlands) and then excised as a ClaI-NheI fragment and inserted into ClaI- and NheI-digested pBP23 [50], a suicide vector containing the *E. coli* ampG gene with flanking *B. pertussis* DNA of 618 bp (from position 50,474 to 51,092 of the *B. pertussis* genome) and 379 bp (from position 52,581 to 52,960 of the *B. pertussis* genome) at the 5' and 3' end of *E. coli* ampG, respectively. The resulting plasmid was

12

transferred into *E. coli* SM10 [51], which was then conjugated with BPSM, and two successive homologous recombination events were selected as described [52]. Ten individual colonies were screened by PCR as follows. The colonies were suspended in 100 μ l H₂O, heated for 20 min. at 95° C., and centrifuged for 5 min at 15,000 \times g. One μ l of supernatants was then used as template for PCR using oligonucleotides A and C: 5'-TAAGAAGCAAATAAGCCAGGCATT-3' (SEQ ID No:7) to verify the presence of *E. coli* ampG and using oligonucleotides D: 5'-TATAC-CATGGCGCCGCTGCTGGTGCTGGGC-3' (SEQ ID No:8) and E: 5'-TATATCTAGACGCTGGCCGTAACCT-TAGCA-3' (SEQ ID No:9) to verify the absence of *B. pertussis* ampG. One of the strains containing *E. coli* ampG and lacking *B. pertussis* ampG was then selected, and the entire ampG locus was sequenced. This strain was then used for further engineering.

The ptx genes were deleted from the chromosome of this strain as described [21] and then replaced by mutated ptx coding inactive PTX. The EcoRI fragment containing the mutated ptx locus from pPT-RE [16] was inserted into the EcoRI site of pJQ200mp18rpsl [53]. The resulting plasmid was integrated into the *B. pertussis* chromosome at the ptx locus by homologous recombination after conjugation via *E. coli* SM10. The ptx locus in the chromosome of the resulting *B. pertussis* strain was sequenced to confirm the presence of the desired mutations. Toxin production was analyzed by immunoblotting using a mix of monoclonal antibodies IB7 [54] specific for subunit S1, and 11E6 [55] specific for subunits S2 and S3 of PTX.

Finally, the dnt gene was deleted from the resulting *B. pertussis* strain as the dnt flanking regions were amplified by PCR using BPSM genomic DNA as a template and oligonucleotides F: 5'-TATAGAATTCGCTCGGTTTCGCTGGT-CAAG G-3' (SEQ ID No:10) and G: 5'-TATATCTAGAGCAATGCCGATTCATCTTA-3' (SEQ ID No: 11) for the dnt upstream region, and H: 5'-TATATCTAGAGCGGCTT TATTGCTTTTCC-3' (SEQ ID No:12) and I: 5'-TATAAAGCTTCTCATGCACGCCG GCTTCTC-3' (SEQ ID No:13) for the dnt downstream region, as primers. The resulting 799-bp and 712-bp DNA fragments were digested with EcoRI/XbaI and XbaI/HindIII, respectively, and linked together using the Fast Link kit (Epicentre Biotechnologies, Madison, Wis.). The ligated fragment was amplified by PCR using oligonucleotides F and I, and the 1505-bp PCR fragment was then inserted into pCR2.1-Topo (Invitrogen), re-isolated from the resulting plasmid as an EcoRI fragment and inserted into the unique EcoRI site of pJQmp200rpsL18. The resulting plasmid was introduced into *B. pertussis* by conjugation via *E. coli* SM10. Successful deletion of the dnt gene by allelic exchange was verified by Southern blot analysis on PvuII-digested *B. pertussis* genomic DNA using the PCR fragment corresponding to the dnt upstream region as a probe. The probe was labeled with digoxigenin (DIG) using the DIG Easy Hyb labeling kit (Roche, Meylan, France). The sizes of the hybridizing bands were determined from the migration distance of the Dig-labeled DNA molecular marker III (Roche). The dnt locus of this final strain, named BPZE1 was sequenced.

Example 3—Analysis of TCT Production

For sensitive quantitation of TCT production, culture supernatants of *B. pertussis* grown to logarithmic phase were collected, subjected to solid phase extraction [15] and

13

derivatized with phenylisothiocyanate (PITC, Pierce). The resulting phenylthiocarbonyl (PTC) derivatives were separated by reversed-phase HPLC using a C8 column (Perkin Elmer) and detected at 254 nm. The amount of *B. pertussis* PTC-TCT in each sample was determined by comparing the peak area and elution time with an identically processed TCT standard.

Example 4—Cell-Adherence Assay

To analyze adherence properties of the *B. pertussis* strains, their attachment rates to the human pulmonary epithelial cell line A549 (ATCC n° CCL-185) and the murine macrophage cell line J774 (ATCC n° TIB-67) were measured as previously described [16].

Example 5—Transmission Electron Microscopy

The single droplet-negative staining procedure was used as described previously [17] with the following modifications, 20 μ l of a suspension at approximately 10^9 bacteria/ml were absorbed for 2 min. onto form forward carbon-coated nickel grids (400 mesh; Electron Microscopy Sciences EMS, Washington, Pa.). After 30 seconds air-drying the grids were stained for 2 minutes with 20 μ l of 2% phosphotungstic acid (pH7; EMS) and examined after air-drying under a transmission electron microscope (Hitachi 7500, Japan) at 60 kvolts and high resolution.

Example 6—Intranasal Infection and Vaccination

3-week and 8-week old female Balb/C were kept under specific pathogen-free conditions, and all experiments were carried out under the guidelines of the Institut Pasteur de Lille animal study board. Mice were intranasally infected with approximately 4×10^6 bacteria in 20 μ l PBS, and kinetics of CFU in the lungs were measured as previously described [18]. For vaccination with aPV (Tetravac; Aventis-Pasteur, France), mice were immunized intraperitoneally (i.p.) with 20% of the human dose and boosted one month later using the same dose.

Example 7—Antibody Determination

Sera were collected, and antibody titers were estimated by enzyme-linked immunosorbent assays (ELISA) as previously described [18].

Example 8—Cytokine Assays

Spleen cells from individual mice were tested at different time points after immunization for in vitro cytokine production in response to heat-killed *B. pertussis* BPSM (10^6 cells/ml), 5.0 μ g/ml PTX (purified from *B. pertussis* BPGR4 [19] as previously described [20] and heat-inactivated at 80° C. for 20 min), 5.0 μ g filamentous hemagglutinin (FHA, purified from *B. pertussis* BPRA [21] as previously described [22]), 5 μ g/ml concanavalin A (Sigma Chemical Co., St. Louis, Mo.) or medium alone as control. Supernatants were removed from triplicate cultures after 72 h incubation at 37° C. and 5% CO₂, and IFN- γ and IL-5 concentrations were determined by immunoassays (BD OptEIA set, Pharmingen).

Example 9—Intranasal Infection and Vaccination: Challenge at 1, 2, 3 and 4 Weeks

An infant (3 weeks-old) mouse model [29] was used to compare the efficiency of vaccination with BPZE1 with the

14

one of vaccination with acellular pertussis vaccine (aPv). Female Balb/C mice were intranasally infected with approximately 1×10^6 BPZE1 strain in 20 μ l PBS. For vaccination with aPv (Tetravac; Aventis-Pasteur, France), mice were immunized intraperitoneally with 20% of the human dose. One, two, three or four weeks after vaccination with BPZE1 or aPv, mice were intranasally challenged with virulent *B. pertussis* BPSM/bctA-lacZ strain [53]. This strain is a BPSM-derivative gentamycin-resistant which allows the discrimination with BPZE1 (gentamycin-sensitive) on Bordet-Gengou agar plates containing 10 μ g/ml of gentamycin and 100 μ g/ml of streptomycin (BGgs). Control group corresponds to naive mice challenged with BPSM/bctA-lacZ. One week after challenge infection, lungs were aseptically removed, homogenized and plates on BGgs for CFU determination as previously described [18].

Mice were vaccinated with BPZE1 or aPv and challenged with virulent *B. pertussis* one, two, three or four weeks after vaccination. Lung CFUs counts were determined 3 hours or 7 days later. Results are expressed as mean (\pm standard error) CFUs from three to five mice per group. Levels of protection are calculated for each challenge infection as mean percentages of CFUs of each group relative of the average of CFUs in non-immunized group, 7 days after challenge infection (Tables 2 to 5).

Example 10—Statistical Analysis

The results were analyzed using the unpaired Student's t test and the Kruskal-Wallis test followed by the Dunn's post-test (GraphPad Prism program) when appropriate. Differences were considered significant at P=0.05.

Results

Construction of *B. pertussis* BPZE1

Three virulence factors were genetically targeted: tracheal cytotoxin (TCT), pertussis toxin (PTX) and dermonecrotic toxin (DNT).

TCT is responsible for the destruction of ciliated cells in the trachea of infected hosts [24, 25] and may thus be involved in the cough syndrome. TCT is a breakdown product of peptidoglycan in the cell wall of Gram-negative bacteria, which generally internalize it into the cytosol by the AmpG transporter protein to be re-utilized during cell wall biosynthesis. *B. pertussis* AmpG is inefficient in the internalization of peptidoglycan breakdown products. We therefore replaced the *B. pertussis* ampG gene by *E. coli* ampG. The resulting strain expressed less than 1% residual TCT activity (FIG. 1).

PTX is a major virulence factor responsible for the systemic effects of *B. pertussis* infections and is composed of an enzymatically active moiety, called S1, and a moiety responsible for binding to target cell receptors (for review, see 26). However, it is also one of the major protective antigens, which has prompted us to replace the natural ptx genes by a mutated version coding for an enzymatically inactive toxin. This was achieved by replacing Arg-9 by Lys and Glu-129 by Gly in S1, two key residues involved in substrate binding and catalysis, respectively. Allelic exchange was used to first delete the ptx operon, and then to insert the mutated version. The presence of the relevant toxin analogues in the *B. pertussis* culture supernatants was evaluated by immunoblot analysis (FIG. 2).

Finally, allelic exchange was used to remove the dnt gene (FIG. 3). Although the role of DNT in the virulence of *B. pertussis* is not certain, it has been identified as an important

toxin in the closely related species *Bordetella bronchiseptica* and displays lethal activity upon injection of minute quantities (for review, see 26).

In Vitro Characterization of *B. pertussis* BPZE1

Since some of the genetic alterations in BPZE1 may potentially affect the bacterial cell wall synthesis, the size and shape, as well as the in vitro growth rate of BPZE1 was compared with those of the parental strain BPSM. The growth rate of BPZE1 did not differ from that of BPSM (FIG. 4), and no difference in bacterial shape or size was detected between BPZE1 and BPSM, as evidenced by electron microscopy analysis (FIG. 5). However, the cell wall of BPZE1 appeared to be consistently somewhat thinner than that of BPSM.

To determine whether the absence or alterations of any of the targeted toxins in BPZE1 affects adherence properties of *B. pertussis*, the attachment rates of BPZE1 was compared with those of BPSM, using the human pulmonary epithelial cell line A549 and the murine macrophage cell line J774, as two cellular models often used to study the adherence of *B. pertussis*. No significant difference in the adherence capacities to either cell line was observed between the two strains (FIG. 6).

Attenuation of *B. pertussis* BPZE1

To determine whether the mutations introduced into *B. pertussis* BPZE1 have resulted in attenuation, yet allow the organism to colonize the respiratory tract, Balb/C mice were intranasally infected with BPZE1 or BPSM, and colonization was followed over time. BPZE1 was able to colonize and persist in the lungs of mice as long as BPSM (FIG. 7). However, the peak of multiplication seen 7 days after infection with BPSM was consistently lacking in mice infected with BPZE1. Studies done with strains mutated in individual toxin genes indicated that this is due to the mutations in the *ptx* locus (data not shown). When the lungs were examined for histopathological changes and inflammatory infiltration, infection with BPSM was found to induce strong peri-bronchiovascular infiltrates and inflammatory cell recruitment 7 days after infection, associated with a strong hypertrophy of the bronchiolar epithelial cells (FIG. 8). In contrast, no such changes were seen in BPZE1-infected animals, and the histology of the BPZE1-infected mice was similar to that of the control mice that had received PBS instead of the bacteria. The BPSM-infection induced inflammation lasted for at least two months (data not shown). These results indicate that the mutations introduced into BPZE1 have resulted in drastic attenuation, but allow the bacteria to colonize and persist in the lungs.

Protection Against *B. pertussis* Challenge after Intranasal Vaccination of Adult Mice with BPZE1

To evaluate the protection offered by BPZE1, the effect of a single intranasal administration of this strain to 8-weeks old Balb/C mice on the subsequent colonization by the wild type challenge strain BPSM was compared with that of two i. p. immunizations with 1/2 of a human dose of aPV. This aPV immunization protocol has been described as the best to correlate with pertussis vaccine efficacy in human clinical trials [27, 28]. As shown by the total clearance of bacterial colony counts in the lungs seven days after challenge infection, a single intranasal administration of BPZE1 and two i.p. immunizations with aPV provided similar levels of protection (FIG. 9a). High bacterial loads were found in the control mice that had received two injections of PBS instead of the vaccine.

Protection Against *B. pertussis* Challenge after Intranasal Vaccination of Infant Mice with BPZE1

Since the principal targets of novel pertussis vaccines are young infants, that are not protected with the currently available vaccines, an infant (3 weeks-old) mouse model [29] was developed and used to compare the efficiency of vaccination with BPZE1 with that of vaccination with aPV. A single nasal administration of BPZE1 fully protected infant mice against challenge infection (FIG. 9b), as complete bacterial clearance was observed in the lungs one week after challenge. In contrast, substantial numbers of bacteria remained in the aPV-vaccinated animals one week after challenge infection. The difference in bacterial load between the BPZE1-vaccinated and the aPV-vaccinated mice was statistically significant, indicating that in the infant mouse model a single intranasal administration with BPZE1 provides better protection than two systemic administrations of aPV.

In addition, a strong reduction in the bacterial load of the challenge strain 3 hours after administration when the mice had been immunized with BPZE1 was consistently observed compared to the aPV-immunized animals (FIG. 9c), indicating that vaccination with BPZE1 reduces the susceptibility to infection by the challenge strain. This effect was seen in both 8-weeks old and in infant mice. In contrast, aPV had no effect on the bacterial counts 3 hours after infection, when compared to the control mice.

Protection Against *B. paraptussis* Challenge after Intranasal Vaccination with BPZE1

There is increasing concern about *B. paraptussis* infection in children, especially in immunized populations [30, 31]. *B. paraptussis* causes a milder pertussis-like syndrome, the frequency of which is probably largely underestimated. Furthermore, the incidence of *B. paraptussis* infections has been increasing over the last decades, possibly due to the fact that pertussis vaccines are known to have very low or no protective efficacy against *B. paraptussis* [32, 33]. In contrast, infection by *B. pertussis* has recently been reported to protect against *B. paraptussis* infection [34]. BPZE1 was also assessed for protection against *B. paraptussis* using the infant mouse model. Whereas two administrations of aPV did not provide any protection against *B. paraptussis*, as previously reported, a single intranasal administration of BPZE1 provided strong protection, as measured by the low numbers of *B. paraptussis* counts in the lungs of the vaccinated mice 1 week after challenge (FIG. 9d).

Immune Responses Induced by BPZE1 Vaccination

Although the mechanisms of protective immunity against *B. pertussis* infection are not yet completely understood, clear evidence of a role for both B cells and IFN- γ has been demonstrated in mice [28]. Vaccination with either one nasal dose of BPZE1 or two i. p. administrations of aPV induced high titers of serum IgG against FHA, a major surface antigen of *B. pertussis* 1351, also present in aPV (FIG. 10a). Following *B. pertussis* challenge, positive anamnestic responses were measured in BPZE1- and in aPV-vaccinated animals, as indicated by an increase in anti-FHA IgG titers, compared to primary responses before *B. pertussis* infection. Examination of the anti-FHA IgG1/IgG2a ratios showed that these ratios were higher after aPV administration, characteristic of a Th2 type response, than after BPZE1 vaccination (FIG. 10b). Although the anti-FHA-IgG/IgG2a decreased after challenge in the aPV vaccinated mice, it remained still substantially higher than in the BPZE1-vaccinated animals after *B. pertussis* challenge.

17

Analysis of *B. pertussis* antigen-specific cytokine patterns induced by BPZE1 or aPv vaccination confirmed that BPZE1 administration favors a stronger Th1 type response than aPv vaccination. This was revealed by the fact that the ratios of IFN- γ over IL-5 produced by splenocytes stimulated with FHA or PT, or with the polyclonal activator ConA were significantly higher in BPZE1 vaccinated mice than in aPv vaccinated mice (FIG. 10c).

Protective Immunity of BPZE1 Over Time (from 1 Week to 4 Weeks)

As shown in Tables 1 to 5 below, whereas administration of aPv provided limited protection (reduction of 75% of bacterial load compared to non-vaccinated mice at 1 week) against *B. pertussis*, a single intranasal administration of BPZE1 already provided high level of protection (reduction of 97.64% of bacterial load) against a *B. pertussis* challenge infection performed one week after vaccination. If challenge infection occurred two weeks after vaccination, the level of protection induced by BPZE1 reached more than 99.999% compared to non-vaccinated mice and is significantly superior to the protection induced by aPv vaccine (approximately 92% compared to non-vaccinated mice). Therefore, vaccine efficacy of BPZE1 against *B. pertussis* challenge is already significant one week after vaccination and is increasing over the at least next two months.

TABLE 1

Kinetics of vaccines efficacy against <i>B. pertussis</i> challenge in infant mice.				
Time between vaccination and challenge	Time between lungs recovery and challenge	Log10 cfu/lungs of mice		
		Naive	aPv-vaccinated	BPZE1-vaccinated
1 week	3 hours	5.71 \pm 0.03	5.8 \pm 0.07	5.74 \pm 0.01
	7 days	6.71 \pm 0.06	5.97 \pm 0.20	4.86 \pm 0.35
2 weeks	3 hours	5.77 \pm 0.10	5.60 \pm 0.02	5.49 \pm 0.05
	7 days	6.49 \pm 0.08	5.31 \pm 0.16	3.22 \pm 0.33
3 weeks	3 hours	6.03 \pm 0.11	5.88 \pm 0.04	5.33 \pm 0.08
	7 days	6.58 \pm 0.09	5.62 \pm 0.11	3.14 \pm 0.38
4 weeks	3 hours	6.31 \pm 0.01	6.15 \pm 0.02	5.83 \pm 0.05
	7 days	6.36 \pm 0.04	5.21 \pm 0.11	1.83 \pm 0.46

TABLE 2

Level of protection of aPv-vaccinated and BPZE1-vaccinated mice as compared to non-vaccinated mice at week 1.				
Non vaccinated mice	Number of bacteria in lungs	Mean number of bacteria		
Non-vaccinated 1	4.7 $\times 10^6$	5.36 $\cdot 10^6$		
Non-vaccinated 2	3.8 $\times 10^6$			
Non-vaccinated 3	8.2 $\times 10^6$			
Non-vaccinated 4	4.1 $\times 10^6$			
Non-vaccinated 5	6 $\times 10^6$			

Non vaccinated mice	Number of bacteria in lungs	Percentage of remaining bacteria ⁽¹⁾	Mean percentage of remaining bacteria	Level of protection
aPv1	1.95 $\times 10^6$	36.38	25%	75%
aPv2	2.9 $\times 10^6$	54.1		
aPv3	2.9 $\times 10^5$	5.41		
aPv4	3.6 $\times 10^5$	6.72		
aPv5	1.2 $\times 10^6$	22.39		

18

TABLE 2-continued

Level of protection of aPv-vaccinated and BPZE1-vaccinated mice as compared to non-vaccinated mice at week 1.				
BPZE1-vaccinated mice				
BPZE1-1	3.2 $\times 10^5$	5.97	2.36%	97.64%
BPZE1-2	2 $\times 10^4$	0.004		
BPZE1-3	6 $\times 10^4$	1.12		

⁽¹⁾ Percentage of remaining bacteria = number of bacteria for each particular mouse/mean number of bacteria of all non-vaccinated mice

TABLE 3

Level of protection of aPv-vaccinated and BPZE1-vaccinated mice as compared to non-vaccinated mice at week 2.				
Non vaccinated mice	Number of bacteria in lungs	Mean number of bacteria		
Non-vaccinated 1	5 $\times 10^6$	3.34 $\times 10^6$		
Non-vaccinated 2	3.6 $\times 10^6$			
Non-vaccinated 3	1.7 $\times 10^6$			
Non-vaccinated 4	2.4 $\times 10^6$			
Non-vaccinated 5	4 $\times 10^6$			

Non vaccinated mice	Number of bacteria in lungs	Percentage of remaining bacteria ⁽¹⁾	Mean percentage of remaining bacteria	Level of protection
aPv1	9.5 $\times 10^4$	2.84	8.11%	91.89%
aPv2	2.9 $\times 10^5$	8.68		
aPv3	1 $\times 10^5$	2.99		
aPv4	6.8 $\times 10^5$	20.36		
aPv5	1.9 $\times 10^5$	5.69		

BPZE1-vaccinated mice				
BPZE1-1	9.5 $\times 10^3$	2.8 $\times 10^{-3}$	1.03 $\times 10^{-3}$ %	99.999%
BPZE1-2	450	1.35 $\times 10^{-4}$		
BPZE1-3	3500	1.05 $\times 10^{-3}$		
BPZE1-4	500	1.5 $\times 10^{-4}$		

⁽¹⁾ Percentage of remaining bacteria = number of bacteria for each particular mouse/mean number of bacteria of all non-vaccinated mice

TABLE 4

Level of protection of aPv-vaccinated and BPZE1-vaccinated mice as compared to non-vaccinated mice at week 3.				
Non vaccinated mice	Number of bacteria in lungs	Mean number of bacteria		
Non-vaccinated 1	1.8 $\times 10^6$	4.04 $\times 10^6$		
Non-vaccinated 2	5.75 $\times 10^6$			
Non-vaccinated 3	4.7 $\times 10^6$			
Non-vaccinated 4	3.2 $\times 10^6$			
Non-vaccinated 5	4.75 $\times 10^6$			

Non vaccinated mice	Number of bacteria in lungs	Percentage of remaining bacteria ⁽¹⁾	Mean percentage of remaining bacteria	Level of protection
aPv1	1.99 $\times 10^5$	4.94	11.26%	88.74%
aPv2	6 $\times 10^5$	14.85		
aPv3	6 $\times 10^5$	14.85		
aPv4	4.2 $\times 10^5$	10.40		

BPZE1-vaccinated mice				
BPZE1-1	3640	9.01 $\times 10^{-4}$	8.65 $\times 10^{-4}$ %	99.999%
BPZE1-2	9720	2.4 $\times 10^{-3}$		

TABLE 4-continued

Level of protection of aPv-vaccinated and BPZE1-vaccinated mice as compared to non-vaccinated mice at week 3.		
BPZE1-3	300	7.43×10^{-5}
BPZE1-4	340	8.42×10^{-5}

⁽¹⁾ Percentage of remaining bacteria = number of bacteria for each particular mouse/mean number of bacteria of all non-vaccinated mice

TABLE 5

Level of protection of aPv-vaccinated and BPZE1-vaccinated mice as compared to non-vaccinated mice at week 4.				
Non vaccinated mice	Number of bacteria in lungs	Mean number of bacteria		
Non-vaccinated 1	2.1×10^6	2.36×10^6		
Non-vaccinated 2	2.2×10^6			
Non-vaccinated 3	3.1×10^6			
Non-vaccinated 4	2.6×10^6			
Non-vaccinated 5	1.8×10^6			
	Number of bacteria in lungs	Percentage of remaining bacteria ⁽¹⁾	Mean percentage of remaining bacteria	Level of protection
aPv-vaccinated mice				
aPv1	2.52×10^5	10.68	7.76%	92.24%
aPv2	3.28×10^5	13.90		
aPv3	1.04×10^5	4.41		
aPv4	8.4×10^5	3.56		
aPv5	1.48×10^5	6.27		
BPZE1-vaccinated mice				
BPZE1-1	190	8.05×10^{-5}	$7.13 \times 10^{-5}\%$	99.999%
BPZE1-2	0	0		
BPZE1-3	110	4.66×10^{-5}		
BPZE1-4	320	1.36×10^{-4}		
BPZE1-5	220	9.32×10^{-5}		

⁽¹⁾ Percentage of remaining bacteria = number of bacteria for each particular mouse/mean number of bacteria of all non-vaccinated mice

DISCUSSION

Pertussis is the first infectious disease whose incidence is increasing in countries with high vaccine coverage. This paradoxical situation is most likely linked to the epidemiological changes observed since the massive introduction of highly efficacious vaccines. In contrast to the pre-vaccination era, cases of adolescent and adult pertussis are now increasingly more frequent. Although generally not life-threatening in that age group, *B. pertussis*-infected adults are an important reservoir for infection of the very young children, too young to be protected by vaccination. Early vaccination, possibly at birth, would therefore be highly desirable, but is hampered by the immaturity of the immune system of neonates and infants. However, the fact that natural *B. pertussis* infection, even very early in life, is able to induce a strong Th1 response in infants [12] prompted us to develop a live attenuated *B. pertussis* vaccine strain to be given by the nasal route as an alternative over the currently available vaccines.

Based on experimental infections of primates, Huang et al. had already in 1962 come to the conclusion that ultimate protection against whooping cough probably best follows a live *B. pertussis* inoculation [36]. In veterinary medicine, attenuated *Bordetella* strains have been used to vaccinate against bordetellosis in dogs and piglets. A live attenuated *Bordetella bronchiseptica* strain has been shown to provide

strong protection against kennel cough in dogs 1371 after nasal administration. This protection was seen as early as 48 h after vaccination. Intranasal vaccination with live attenuated *B. bronchiseptica* has also been shown to protect against atrophic rhinitis in two-days old piglets [38], indicating that in a live attenuated form *Bordetella* vaccines can be highly active in new-born animals.

Previous attempts to genetically attenuate *B. pertussis* as a live vaccine candidate have met with rather limited success. Based on a strategy used for the successful attenuation of *Salmonella* vaccine strains [39], Roberts et al. have deleted the *aroA* gene of *B. pertussis* 1401. The *aroA* mutant was indeed highly attenuated, but had also lost its capacity to colonize the respiratory tract of the intranasally vaccinated animals and induced protective immunity only after repeated administrations of high doses. We took advantage of the knowledge on the molecular mechanisms of *B. pertussis* virulence and developed the highly attenuated strain BPZE1. This strain contains genetic alterations leading to the absence or inactivation of three major toxins, PTX, TCT and DNT. In contrast to the *aroA* mutant, this strain was able to colonize the mouse respiratory tract and to provide full protection after a single intranasal administration. The protection in adult mice was indistinguishable from that induced by two administrations of $\frac{1}{5}$ of a human dose of aPV. An important difference, however, was seen in infant mice, where a single administration of BPZE1 fully protected, whereas aPV only offered partial protection. In the context of the difficulties to induce protection in infants with the currently available vaccines early in life, these results provide hope for the development of novel vaccination strategies that may be used in the very young children, possibly at birth. In addition, BPZE1 protected against *B. parapertussis*, whereas aPV did not. Therefore the use of BPZE1 should also have an impact on the incidence of whooping cough caused by *B. parapertussis* in infants.

Although the recent replacement of first generation whole-cell vaccines by new aPV in many countries has significantly reduced the systemic adverse reactions observed with whole-cell vaccines, it has not abolished the need for repeated vaccination to achieve protection. This makes it unlikely to obtain protection in very young children (<6 months) that present the highest risk to develop severe disease. In addition, the wide-spread use of aPV has revealed new, unforeseen problems. Repeated administration of aPV may cause extensive swelling at the site of injection [41], which was not observed with whole-cell vaccines. In approximately 5% of the cases this swelling involves almost the entire limb and lasts for more than a week. Although the mechanism of this swelling has not been characterized yet, it has been proposed to be due to an Arthus hypersensitivity reaction caused by high antibody levels induced by the primary immunization [42]. However, it could also be related to the Th2 skewing of the immune response, as, compared to whole-cell vaccines, aPV administration induces more Th2-type cytokines in vaccinated children [10] and causes a delay in the Th1 development (Mascart et al., in preparation). Delayed maturation of Th1 function has been associated with a risk for atopy in genetically predisposed individuals [133]. The two mechanisms are not mutually exclusive. Compared to aPV, the immune response to BPZE1 administration is less biased towards the Th2 arm, and since BPZE1 is administered mucosally, no swelling reaction can occur.

The use of live attenuated bacteria as vaccines raises the issue of their biosafety. As such, they fall under the directives and guidelines for genetically modified organisms

susceptible to be released into the environment. These guidelines and directives describe several objectives that have to be met, including hazard identification and environmental risk assessment [44]. Potential pathogenicity needs to be carefully considered, especially in immuno-compromised individuals, such as those infected with HIV. The natural biology of *B. pertussis* is particularly interesting in that regard. Although pertussis in HIV-infected subjects has been described occasionally, it is rather rare in AIDS patients [45]. In its genetically attenuated form, *B. pertussis* would therefore not be expected to cause major problems in HIV-infected children, especially if severe AIDS is an exclusion criterion, as it is for many vaccines. *B. pertussis* colonization is strictly limited to the respiratory epithelium, without extrapulmonary dissemination of the bacteria, which naturally excludes systemic bacteremia of the BPZE1 vaccine strain. If nevertheless unforeseeable safety problems occurred, the vaccine strain can easily be eliminated by the use of macrolide antibiotics, such as erythromycin, to which essentially all *B. pertussis* isolates are highly sensitive.

A further concern, like for any live vaccine, is the potential release of the vaccine strain in the environment and the consequences of such a release. *B. pertussis* is a strictly human pathogen, and there is no animal vector or reservoir. Moreover, unlike *B. bronchiseptica*, survival of wild-type *B. pertussis* in the environment is extremely limited [46]. Pertussis is only spread by coughing individuals, and there appears to be no asymptomatic carriage [47]. Coughing cannot be assessed in the mouse models used in this study. However, due to the nature of the genetic alterations in BPZE1, in particular the strong reduction of TCT and the genetic inactivation of PTX, this strain would not be expected to induce coughing. Active PTX has been shown to be required for cough induction in a coughing rat model, although the mechanism is not known [48]. If the vaccine strain were nevertheless to be transmitted to non-vaccinated individuals, this would at worst result in increased vaccine coverage. The consequences of each of these potential hazards can thus be graded as negligible and can easily and rapidly be controlled by antibiotic treatment if necessary.

Advantages of the use of BPZE1 include the relatively low production costs, making it especially attractive for developing countries, its needle-free easy and safe mode of administration and its potential to induce mucosal immunity in addition to systemic immunity. Although the role of mucosal immunity against pertussis has surprisingly not been much addressed, the fact that *B. pertussis* is a strictly mucosal pathogen, makes it likely that mucosal immune responses may contribute significantly to protection. None of the currently available vaccines induces any significant mucosal response.

Other advantages of the use of BPZE1 in vaccination are: the rapid protective immune response obtained after a single intranasal dose of BPZE1, since induction of the immunity can be detected 1 week after vaccination, an increase of the protective immunity over the at least next two months after vaccination, and the complete protective immunity, since a level of protection of more than 99.999% is obtained 2 weeks after vaccination.

The use of live attenuated *B. pertussis* for mucosal vaccination offers yet another advantage. *B. pertussis* can be used for the presentation of heterologous antigens to the respiratory mucosa (for review see 49). The use of BPZE1 as a heterologous expression platform may thus be helpful for the generation of multivalent vaccines against a variety of respiratory pathogens. However, since intranasal delivery

of BPZE1 also induces strong systemic immune responses, as shown here by both the high levels of anti-FHA antibodies and of antigen-specific IFN- γ production, it may also be used for the production of antigens to which systemic immune responses are desired.

While the invention has been described in terms of various preferred embodiments, the skilled artisan will appreciate that various modifications, substitutions, omissions and changes may be made without departing from the scope thereof. Accordingly, it is intended that the scope of the present invention be limited by the scope of the following claims, including equivalents thereof.

REFERENCES

1. WHO (2004) The world health report 2004-changing history, Geneva, WHO.
2. Das P (2002) Whooping cough makes global comeback. *Lancet* ii: 322.
3. Tan T, Trindade E, Skowronski D (2005) Epidemiology of Pertussis. *Pediatr Infect Dis J* 24: S10-S18.
4. Centers for Disease Control and Prevention. Epidemiology and Prevention of Vaccine-Preventable Diseases. Atkinson W, Wolfe S, Hamborsky J, McIntyre L, eds. 9th ed. Washington D.C.: Public Health Foundation, 2006; Pertussis Chapter 14.
5. Wirsing von König C H, Halperin S, Riffelmann M, Guiso N (2002) Pertussis of adults and infants. *Lancet Infect Dis* 2: 744-750.
6. Lewis D B, Yu C C, Meyer J, English B K, Kahn S J, et al. (1991) Cellular and molecular mechanisms for reduced interleukin-4 and interferon- γ production by neonatal T cells. *J Clin Invest* 87: 194-202.
7. Siegrist C A (2001) Neonatal and early life vaccinology. *Vaccine*. 19: 3331-3346.
8. Mills K H G (2001) Immunity to *Bordetella pertussis*. *Microbes Infect* 3: 655-677.
9. Lewis D B, Larsen A, Wilson C B (1986) Reduced interferon- γ mRNA levels in human neonates. *J Exp Med* 163:1018-1023.
10. Ausiello C M, Urbani F, La Sala A, Lande R, Cassone A (1997) Vaccine- and antigen-dependent type 1 and type 2 cytokine induction after primary vaccination in infants with whole-cell or acellular pertussis vaccines. *Infect Immun* 65: 2168-2174.
11. Wirsing von König C H, Postels-Multani S, Bock H L, Schmitt H J (1995) Pertussis in adults: frequency of transmission after household exposure. *Lancet* 346:1326-1329.
12. Mascart F, Verscheure V, Malfroot A, Hainaut M, Piérard D, et al. (2003) *Bordetella pertussis* infection in 2-months-old infants promotes Type 1 T cell responses. *J Immunol* 170:1504-1509.
13. Menozzi F D, Mutombo R, Renauld G, Gantiez C, Hannah J H, et al. (1994) Heparin-inhibitable lectin activity of the filamentous hemagglutinin adhesin of *Bordetella pertussis*. *Infect Immun* 62: 769-778.
14. Imaizumi A, Suzuki Y, Ono S, Sato H, Sato Y (1983) Effect of heptakis (2,6-O-dimethyl)-beta-cyclodextrin on the production of pertussis toxin by *Bordetella pertussis*. *Infect Immun* 41: 1138-1143.
15. Cookson B T, Cho H-L, Herwaldt L A, Goldman W E (1989) Biological activities and chemical composition of purified tracheal cytotoxin of *Bordetella pertussis*. *Infect Immun* 57: 2223-2229.
16. Alonso S, Pethe K, Mielcarek N, Raze D, Loch C (2001) Role of ADP-ribosyltransferase activity of pertussis toxin

- in toxin-adhesin redundancy with filamentous hemagglutinin during *Bordetella pertussis* infection. *Infect Immun* 69: 6038-6043.
17. Collyn F, Lety M A, Nair S, Escuyer V, Ben Younes A, et al. (2002) *Yersinia pseudotuberculosis* harbors a type IV pilus gene cluster that contributes to pathogenicity. *Infect Immun* 70: 619-620.
 18. Mielcarek N, Comette J, Schacht A M, Pierce R J, Locht C, et al. (1997) Intranasal priming with recombinant *Bordetella pertussis* for the induction of a systemic immune response against a heterologous antigen. *Infect Immun* 65: 544-550.
 19. Locht C, Geoffroy M C, Renaud G (1992) Common accessory genes for the *Bordetella pertussis* filamentous hemagglutinin and fimbriae share sequence similarities with the papC and papD gene families. *EMBO J* 11: 3175-3183.
 20. Sekura R D, Fish F, Manclark C R, Meade B, Zhang Y L (1983) Pertussis toxin. Affinity purification of a new ADP-ribosyltransferase. *J Biol Chem* 258: 14647-14651.
 21. Antoine R, Locht C (1990) Roles of the disulfide bond and the carboxy-terminal region of the S1 subunit in the assembly and biosynthesis of pertussis toxin. *Infect Immun* 58:1518-1526.
 22. Menozzi F O, Gantiez C, Locht C (1991) Interaction of the *Bordetella pertussis* filamentous haemagglutinin with heparin. *FEMS Microbiol Lett* 62: 59-64.
 23. Locht C, Antoine R, Jacob-Dubuisson F (2001) *Bordetella pertussis*, molecular pathogenesis under multiple aspects. *Curr Opin Microbiol* 4: 82-89.
 24. Heiss L N, Flak T A, Lancaster J R, McDaniel M L, Goldman W E (1993) Nitric oxide mediates *Bordetella pertussis* tracheal cytotoxin damage to the respiratory epithelium. *Infect Agents Dis* 2: 173-177.
 25. Goldman W E, Cookson B T (1988) Structure and functions of the *Bordetella* tracheal cytotoxin. *Tokai J Exp Clin Med* 13 Suppl: 187-191.
 26. Locht C, Antoine R (1999) *Bordetella pertussis* protein toxins. In: Alouf J E, Freer J H, editors. *Comprehensive sourcebook of bacterial protein toxins*. Academic Press, pp. 130-146.
 27. Guiso N, Capiou C, Carletti G, Poolman J, Hauser P (1999) Intranasal murine model of *Bordetella pertussis* infection. I. Prediction of protection in human infants by acellular vaccines. *Vaccine* 17: 2366-2376.
 28. Mills K H, Ryan M, Ryan E, Mahon B P (1998) A murine model in which protection correlates with pertussis vaccine efficacy in children reveals complementary roles for humoral and cell-mediated immunity in protection against *Bordetella pertussis*. *Infect Immun* 66: 594-602.
 29. Roduit C, Bozzotti P, Mielcarek N, Lambert P H, Del Giudice G, et al. (2002) Immunogenicity and protective efficacy of neonatal immunization against *Bordetella pertussis* in a murine model: Evidence for early control of Pertussis. *Infect Immun* 70: 3521-3528.
 30. He Q, Viljanen M K, Arvilommi H, Aittanen B, Mertsola J (1998) Whooping cough caused by *Bordetella pertussis* and *Bordetella parapertussis* in an immunized population. *JAMA* 280: 635-637.
 31. Watanabe M, Nagai M (2004) Whooping cough due to *Bordetella parapertussis*: an unresolved problem. *Expert Rev Anti Infect Ther* 2: 447-454.
 32. Mastrantonio P, Stefanelli P, Giuliano M, Herrera Rojas Y, Ciofi degli Atti M, et al. (1998) *Bordetella parapertussis* infection in children: epidemiology, clinical symptoms, and molecular characteristics of isolates. *J Clin Microbiol* 36: 999-1002.

33. Liese J G, Renner C, Stojanov S, Belohradsky B H, Munich Vaccine Study Group. (2003) Clinical and epidemiological picture of *B. pertussis* and *B. parapertussis* infections after introduction of acellular pertussis vaccines. *Arch Dis Child* 88: 684-687.
34. Watanabe M, Nagai M (2001) Reciprocal protective immunity against *Bordetella pertussis* and *Bordetella parapertussis* in a murine model of respiratory infection. *Infect Immun* 69: 6981-6986.
35. Locht C, Bertin P, Menozzi F D, Renaud G (1993) The filamentous haemagglutinin, a multifaceted adhesin produced by virulent *Bordetella* spp. *Mol Microbiol* 9: 653-660.
36. Huang C C, Chen P M, Kuo J K, Chui W H, Lin S T, et al. (1962) Experimental whooping cough. *N Engl J Med* 266: 105-111.
37. Bey R F, Shade F J, Goodnow R A, Johnson R C (1981) Intranasal vaccination of dogs with live avirulent *Bordetella bronchiseptica*: correlation of serum agglutination titer and the formation of secretory IgA with protection against experimentally induced infectious tracheobronchitis. *Am J Vet Res* 42: 1130-1132.
38. De Jong M F (1987) Prevention of atrophic rhinitis in piglets by means of intranasal administration of a live non-AR-pathogenic *Bordetella bronchiseptica* vaccine. *Vet Q* 9:123-133.
39. Hoiseth S K, Stocker B A D (1981) Aromatic-dependent *Salmonella typhimurium* are non-virulent and effective as live vaccines. *Nature* 291: 238-239.
40. Roberts M, Maskell D, Novotny P, Dougan G (1990) Construction and characterization in vivo of *Bordetella pertussis* aroA mutants. *Infect Immun* 58: 732-739.
41. Rennels M B (2003) Extensive swelling reactions occurring after booster doses of diphtheria-tetanus-acellular pertussis vaccines. *Semin Pediatr Infect Dis* 14: 196-198.
42. Robbins J B, Schneerson R, Trollfors B, Sato H, Sato Y, et al. (2005) The diphtheria and pertussis components of diphtheria-tetanus toxoids-pertussis vaccine should be genetically inactivated mutant toxins. *J Infect Dis* 191: 81-88.
43. Holt P G, Clough J B, Holt B J, Baron-Hay M J U, Rose A H, et al. (1992) Genetic "risk" for atopy is associated with delayed postnatal maturation of T-cell competence. *Clin Exp Allergy* 22: 1093-1099.
44. Favre D, Viret J F (2006) Biosafety evaluation of recombinant live oral bacterial vaccines in the context of European regulation. *Vaccine* May 1; 24 (18):3856-64.
45. Cohn S E, Knorr K L, Gilligan P H, Smiley M L, Weber D J (1993) Pertussis is rare in human immunodeficiency virus disease. *Am Rev Respir Dis* 147: 411-413.
46. Porter J F, Wardlaw A C (1993) Long-term survival of *Bordetella bronchiseptica* in lakewater and in buffered saline without added nutrients. *FEMS Microbiol Lett* 110: 33-36.
47. Linnemann C C Jr, Bass J W, Smith M H D (1968) The carrier state in pertussis. *Am J Epidemiol* 88: 422-427.
48. Parton R, Hall E, Wardlaw A C (1994) Responses to *Bordetella pertussis* mutant strains and to vaccination in the coughing rat model of pertussis. *J Med Microbiol* 40: 307-312.
49. Mielcarek N, Alonso S, Locht C (2001) Nasal vaccination using live bacterial vectors. *Adv Drug Del Rev* 51: 55-69.
50. Lyon R S, Engle J T, Goldman W E. Manuscript in preparation

51. Simon R, Priefer U, Pühler A (1983) A broad host range mobilization system for in vivo genetic engineering: transposon mutagenesis in Gram-negative bacteria. *Bio/Technology* 1: 784-791.
52. Stibitz S (1994) Use of conditionally counterselectable suicide vectors for allelic exchange. *Methods Enzymol* 235: 458-465.
53. Antoine R, Huvent I, Chemial K, Deray I, Raze D, et al. (2005) The periplasmic binding protein of tripartite tri-carboxylate transporter is involved in signal transduction. *J Mol Biol* 351: 799-809.
54. Sato H, Ito A, Chiba J, Sato Y (1984) Monoclonal antibodies against pertussis toxin: effect on toxin activity and pertussis infections. *Infect Immun* 46: 422-428.
55. Sato H, Sato Y, Ito A, Ohishi I (1987) Effect of monoclonal antibody to pertussis toxin on toxin activity. *Infect Immun* 55: 909-915.
56. Tuomanen, E. And Weiss A. (1985) Characterization of two adhesions of *Bordetella pertussis* for human ciliated respiratory epithelial cells. *J. Infect. Dis.* 152:118-125.
57. Locht, C., Antoine, R., Veithen A. and Raze D. 2000. Pertussis Toxin: Structure-Function-Relationship. In Aktories K. Just I editors. Handbook of Experimental Pharmacology, Bacterial Protein Toxins. Springer, vol 145. pp. 167-185.
58. Horiguchi Y, Matsuda, H. Koyama H, Nakai T and Kume K. (1992) *Bordetella bronchiseptica* dermonecrotizing toxin suppresses in vivo antibody responses in mice. *FEMS Microbiol. Lett.* 69:229-234.
59. Bordet et Genysa (1909) L'endotoxine coquelucheuse; *Ann. Inst. Pasteur* 23: 415-419.

60. Iida & Okonogi (1971) Lipo toxicity of *Bordetella pertussis* in mice; *J. Med. Microbiol.* 4: 51-61.
61. R. Parton (1985) Effect of prednisone on the toxicity of *Bordetella pertussis* in mice, *J. Med. Microbiol.* 19: 391-400.
62. Magyar et al (1988) The pathogenesis of turbinate atrophy in pigs caused by *Bordetella bronchiseptica*. *Vet. Microbiol.* 3: 1719-1728.
63. Roop et al (1987) Virulence factors of *Bordetella bronchiseptica* associated with the production of infectious atrophic rhinitis and pneumonia in experimentally infected neonatal swine, *Infect. Immun.* 55: 217-222.
64. Weiss & Goodman (1989) Lethal infection by *Bordetella pertussis* mutants in the infant mouse model, *Infect. Immun.* 57: 3757-3764.
65. Allan & Maskell (1996) The identification, cloning and mutagenesis of a genetic locus required for lipopolysaccharide biosynthesis in *Bordetella pertussis*. *Mol. Microbiol.* 19: 37-52.
66. Alonso et al (2002) Eighty kilodalton N-terminal moiety of *Bordetella pertussis* filamentous hemagglutinin: adherence, immunogenicity, and protective role, *Infection & Immunity.* 70, 4142-4147.
67. Cummings, C. A., Bootsma, H. J., Relman D. A. and Miller J. F. (2006) Species- and Strain-specific Control of a Complex, Flexible Regulon by *Bordetella* BvgAS. *J. Bacteriol.* 188:1775-1785.
68. Kashimoto T., Katahira J, Cornejo W R, Masuda M, Fukuoh A, Matsuzawa T, Ohnishi T, Horiguchi Y. (1999) Identification of functional domains of *Bordetella* dermonecrotizing toxin. *Infect. Immun.* 67 (8) 3727-32.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 13

<210> SEQ ID NO 1

<211> LENGTH: 269

<212> TYPE: PRT

<213> ORGANISM: *Bordetella pertussis*

<400> SEQUENCE: 1

```

Met Arg Cys Thr Arg Ala Ile Arg Gln Thr Ala Arg Thr Gly Trp Leu
 1           5           10           15

Thr Trp Leu Ala Ile Leu Ala Val Thr Ala Pro Val Thr Ser Pro Ala
 20           25           30

Trp Ala Asp Asp Pro Pro Ala Thr Val Tyr Arg Tyr Asp Ser Arg Pro
 35           40           45

Pro Glu Asp Val Phe Gln Asn Gly Phe Thr Ala Trp Gly Asn Asn Asp
 50           55           60

Asn Val Leu Asp His Leu Thr Gly Arg Ser Cys Gln Val Gly Ser Ser
 65           70           75           80

Asn Ser Ala Phe Val Ser Thr Ser Ser Ser Arg Arg Tyr Thr Glu Val
 85           90           95

Tyr Leu Glu His Arg Met Gln Glu Ala Val Glu Ala Glu Arg Ala Gly
 100          105          110

Arg Gly Thr Gly His Phe Ile Gly Tyr Ile Tyr Glu Val Arg Ala Asp
 115          120          125

Asn Asn Phe Tyr Gly Ala Ala Ser Ser Tyr Phe Glu Tyr Val Asp Thr
 130          135          140

Tyr Gly Asp Asn Ala Gly Arg Ile Leu Ala Gly Ala Leu Ala Thr Tyr
 145          150          155          160

```

-continued

Gln Ser Glu Tyr Leu Ala His Arg Arg Ile Pro Pro Glu Asn Ile Arg
 165 170 175
 Arg Val Thr Arg Val Tyr His Asn Gly Ile Thr Gly Glu Thr Thr Thr
 180 185 190
 Thr Glu Tyr Ser Asn Ala Arg Tyr Val Ser Gln Gln Thr Arg Ala Asn
 195 200 205
 Pro Asn Pro Tyr Thr Ser Arg Arg Ser Val Ala Ser Ile Val Gly Thr
 210 215 220
 Leu Val Arg Met Ala Pro Val Ile Gly Ala Cys Met Ala Arg Gln Ala
 225 230 235 240
 Glu Ser Ser Glu Ala Met Ala Ala Trp Ser Glu Arg Ala Gly Glu Ala
 245 250 255
 Met Val Leu Val Tyr Tyr Glu Ser Ile Ala Tyr Ser Phe
 260 265

<210> SEQ ID NO 2
 <211> LENGTH: 1464
 <212> TYPE: PRT
 <213> ORGANISM: Bordetella pertussis

<400> SEQUENCE: 2

Met Asp Lys Asp Glu Ser Ala Leu Arg Gln Leu Val Asp Met Ala Leu
 1 5 10 15
 Val Gly Tyr Asp Gly Val Val Glu Glu Leu Leu Ala Leu Pro Ser Glu
 20 25 30
 Glu Ser Gly Asp Leu Ala Gly Gly Arg Ala Lys Arg Glu Lys Ala Glu
 35 40 45
 Phe Ala Leu Phe Ser Glu Ala Pro Asn Gly Asp Glu Pro Ile Gly Gln
 50 55 60
 Asp Ala Arg Thr Trp Phe Tyr Phe Pro Lys Tyr Arg Pro Val Ala Val
 65 70 75 80
 Ser Asn Leu Lys Lys Met Gln Val Ala Ile Arg Ala Arg Leu Glu Pro
 85 90 95
 Glu Ser Leu Ile Leu Gln Trp Leu Ile Ala Leu Asp Val Tyr Leu Gly
 100 105 110
 Val Leu Ile Ala Ala Leu Ser Arg Thr Val Ile Ser Asp Leu Val Phe
 115 120 125
 Glu Tyr Val Lys Ala Arg Tyr Glu Ile Tyr Tyr Leu Leu Asn Arg Val
 130 135 140
 Pro His Pro Leu Ala Thr Ala Tyr Leu Lys Arg Arg Arg Gln Arg Pro
 145 150 155 160
 Val Asp Arg Ser Gly Arg Leu Gly Ser Val Phe Glu His Pro Leu Trp
 165 170 175
 Phe Ala Tyr Asp Glu Leu Ala Gly Thr Val Asp Leu Asp Ala Asp Ile
 180 185 190
 Tyr Glu Gln Ala Leu Ala Glu Ser Ile Glu Arg Arg Met Asp Gly Glu
 195 200 205
 Pro Asp Asp Gly Ser Leu Asp Thr Ala Glu His Asp Val Trp Arg Leu
 210 215 220
 Cys Arg Asp Gly Ile Asn Arg Gly Glu Gln Ala Ile Phe Gln Ala Ser
 225 230 235 240
 Gly Pro Tyr Gly Val Val Ala Asp Ala Gly Tyr Met Arg Thr Val Ala
 245 250 255
 Asp Leu Ala Tyr Ala Asp Ala Leu Ala Asp Cys Leu His Ala Gln Leu

-continued

260				265				270							
Arg	Ile	Arg	Ala	Gln	Gly	Ser	Val	Asp	Ser	Pro	Gly	Asp	Glu	Met	Pro
		275					280						285		
Arg	Lys	Leu	Asp	Ala	Trp	Glu	Ile	Ala	Lys	Phe	His	Leu	Ala	Ala	Thr
	290					295					300				
Gln	Gln	Ala	Arg	Val	Asp	Leu	Leu	Glu	Ala	Ala	Phe	Ala	Leu	Asp	Tyr
	305				310					315					320
Ala	Ala	Leu	Arg	Asp	Val	Arg	Val	Tyr	Gly	Asp	Tyr	Arg	Asn	Ala	Leu
			325						330					335	
Ala	Leu	Arg	Phe	Ile	Lys	Arg	Glu	Ala	Leu	Arg	Leu	Leu	Gly	Ala	Arg
		340							345				350		
Arg	Gly	Asn	Ala	Ser	Thr	Met	Pro	Ala	Val	Ala	Ala	Gly	Glu	Tyr	Asp
		355					360						365		
Glu	Ile	Val	Ala	Ser	Gly	Ala	Ala	Asn	Asp	Ala	Ala	Tyr	Val	Ser	Met
	370					375							380		
Ala	Ala	Ala	Leu	Ile	Ala	Gly	Val	Leu	Cys	Asp	Leu	Glu	Ser	Ala	Gln
	385				390					395					400
Arg	Thr	Leu	Pro	Val	Val	Leu	Ala	Arg	Phe	Arg	Pro	Leu	Gly	Val	Leu
			405						410					415	
Ala	Arg	Phe	Arg	Arg	Leu	Glu	Gln	Glu	Thr	Ala	Gly	Met	Leu	Leu	Gly
		420							425				430		
Asp	Gln	Glu	Pro	Glu	Pro	Arg	Gly	Phe	Ile	Ser	Phe	Thr	Asp	Phe	Arg
		435					440						445		
Asp	Ser	Asp	Ala	Phe	Ala	Ser	Tyr	Ala	Glu	Tyr	Ala	Ala	Gln	Phe	Asn
	450					455					460				
Asp	Tyr	Ile	Asp	Gln	Tyr	Ser	Ile	Leu	Glu	Ala	Gln	Arg	Leu	Ala	Arg
	465				470					475					480
Ile	Leu	Ala	Leu	Gly	Ser	Arg	Met	Thr	Val	Asp	Gln	Trp	Cys	Leu	Pro
			485						490					495	
Leu	Gln	Lys	Val	Arg	His	Tyr	Lys	Val	Leu	Thr	Ser	Gln	Pro	Gly	Leu
		500							505				510		
Ile	Ala	Arg	Gly	Ile	Glu	Asn	His	Asn	Arg	Gly	Ile	Glu	Tyr	Cys	Leu
		515					520						525		
Gly	Arg	Pro	Pro	Leu	Thr	Asp	Leu	Pro	Gly	Leu	Phe	Thr	Met	Phe	Gln
	530					535					540				
Leu	His	Asp	Ser	Ser	Trp	Leu	Leu	Val	Ser	Asn	Ile	Asn	Gly	Glu	Leu
	545				550					555					560
Trp	Ser	Asp	Val	Leu	Ala	Asn	Ala	Glu	Val	Met	Gln	Asn	Pro	Thr	Leu
			565						570					575	
Ala	Ala	Leu	Ala	Glu	Pro	Gln	Gly	Arg	Phe	Arg	Thr	Gly	Arg	Arg	Thr
		580							585				590		
Gly	Gly	Trp	Phe	Leu	Gly	Gly	Pro	Ala	Thr	Glu	Gly	Pro	Ser	Leu	Arg
		595					600						605		
Asp	Asn	Tyr	Leu	Leu	Lys	Leu	Arg	Gln	Ser	Asn	Pro	Gly	Leu	Asp	Val
	610					615					620				
Lys	Lys	Cys	Trp	Tyr	Phe	Gly	Tyr	Arg	Gln	Glu	Tyr	Arg	Leu	Pro	Ala
	625				630					635					640
Gly	Ala	Leu	Gly	Val	Pro	Leu	Phe	Ala	Val	Ser	Val	Ala	Leu	Arg	His
			645						650					655	
Ser	Leu	Asp	Asp	Leu	Ala	Ala	His	Ala	Lys	Ser	Ala	Leu	Tyr	Lys	Pro
		660							665				670		
Ser	Glu	Trp	Gln	Lys	Phe	Ala	Phe	Trp	Ile	Val	Pro	Phe	Tyr	Arg	Glu
		675					680						685		

-continued

Ile Phe Phe Ser Thr Gln Asp Arg Ser Tyr Arg Val Asp Val Gly Ser
 690 695 700

Ile Val Phe Asp Ser Ile Ser Leu Leu Ala Ser Val Phe Ser Ile Gly
 705 710 715 720

Gly Lys Leu Gly Ser Phe Thr Arg Thr Gln Tyr Gly Asn Leu Arg Asn
 725 730 735

Phe Val Val Arg Gln Arg Ile Ala Gly Leu Ser Gly Gln Arg Leu Trp
 740 745 750

Arg Ser Val Leu Lys Glu Leu Pro Ala Leu Ile Gly Ala Ser Gly Leu
 755 760 765

Arg Leu Ser Arg Ser Leu Leu Val Asp Leu Tyr Glu Ile Phe Glu Pro
 770 775 780

Val Pro Ile Arg Arg Leu Val Ala Gly Phe Val Ser Ala Thr Thr Val
 785 790 795 800

Gly Gly Arg Asn Gln Ala Phe Leu Arg Gln Ala Phe Ser Ala Ala Ser
 805 810 815

Ser Ser Ala Gly Arg Thr Gly Gly Gln Leu Ala Ser Glu Trp Arg Met
 820 825 830

Ala Gly Val Asp Ala Thr Gly Leu Val Glu Ser Thr Ser Gly Gly Arg
 835 840 845

Phe Glu Gly Ile Tyr Thr Arg Gly Leu Gly Pro Leu Ser Glu Cys Thr
 850 855 860

Glu His Phe Ile Val Glu Ser Gly Asn Ala Tyr Arg Val Ile Trp Asp
 865 870 875 880

Ala Tyr Thr His Gly Trp Arg Val Val Asn Gly Arg Leu Pro Pro Arg
 885 890 895

Leu Thr Tyr Thr Val Pro Val Arg Leu Asn Gly Gln Gly His Trp Glu
 900 905 910

Thr His Leu Asp Val Pro Gly Arg Gly Gly Ala Pro Glu Ile Phe Gly
 915 920 925

Arg Ile Arg Thr Arg Asn Leu Val Ala Leu Ala Ala Glu Gln Ala Ala
 930 935 940

Pro Met Arg Arg Leu Leu Asn Gln Ala Arg Arg Val Ala Leu Arg His
 945 950 955 960

Ile Asp Thr Cys Arg Ser Arg Leu Ala Leu Pro Arg Ala Glu Ser Asp
 965 970 975

Met Asp Ala Ala Ile Arg Ile Phe Phe Gly Glu Pro Asp Ala Gly Leu
 980 985 990

Arg Gln Arg Ile Gly Arg Arg Leu Gln Glu Val Arg Ala Tyr Ile Gly
 995 1000 1005

Asp Leu Ser Pro Val Asn Asp Val Leu Tyr Arg Ala Gly Tyr Asp
 1010 1015 1020

Leu Asp Asp Val Ala Thr Leu Phe Asn Ala Val Asp Arg Asn Thr
 1025 1030 1035

Ser Leu Gly Arg Gln Ala Arg Met Glu Leu Tyr Leu Asp Ala Ile
 1040 1045 1050

Val Asp Leu His Ala Arg Leu Gly Tyr Glu Asn Ala Arg Phe Val
 1055 1060 1065

Asp Leu Met Ala Phe His Leu Leu Ser Leu Gly His Ala Ala Thr
 1070 1075 1080

Ala Ser Glu Val Val Glu Ala Val Ser Pro Arg Leu Leu Gly Asn
 1085 1090 1095

-continued

Val Phe Asp Ile Ser Asn Val Ala Gln Leu Glu Arg Gly Ile Gly 1100 1105 1110	Asn Pro Ala Ser Thr Gly Leu Phe Val Met Leu Gly Ala Tyr Ser 1115 1120 1125	Glu Ser Ser Pro Ala Ile Phe Gln Ser Phe Val Asn Asp Ile Phe 1130 1135 1140	Pro Ala Trp Arg Gln Ala Ser Gly Gly Gly Pro Leu Val Trp Asn 1145 1150 1155	Phe Gly Pro Ala Ala Ile Ser Pro Thr Arg Leu Asp Tyr Ala Asn 1160 1165 1170	Thr Asp Ile Gly Leu Leu Asn His Gly Asp Ile Ser Pro Leu Arg 1175 1180 1185	Ala Arg Pro Pro Leu Gly Gly Arg Arg Asp Ile Asp Leu Pro Pro 1190 1195 1200	Gly Leu Asp Ile Ser Phe Val Arg Tyr Asp Arg Pro Val Arg Met 1205 1210 1215	Ser Ala Pro Arg Ala Leu Asp Ala Ser Val Phe Arg Pro Val Asp 1220 1225 1230	Gly Pro Val His Gly Tyr Ile Gln Ser Trp Thr Gly Ala Glu Ile 1235 1240 1245	Glu Tyr Ala Tyr Gly Ala Pro Ala Ala Ala Arg Glu Val Met Leu 1250 1255 1260	Thr Asp Asn Val Arg Ile Ile Ser Ile Glu Asn Gly Asp Glu Gly 1265 1270 1275	Ala Ile Gly Val Arg Val Arg Leu Asp Thr Val Pro Val Ala Thr 1280 1285 1290	Pro Leu Ile Leu Thr Gly Gly Ser Leu Ser Gly Cys Thr Thr Met 1295 1300 1305	Val Gly Val Lys Glu Gly Tyr Leu Ala Phe Tyr His Thr Gly Lys 1310 1315 1320	Ser Thr Glu Leu Gly Asp Trp Ala Thr Ala Arg Glu Gly Val Gln 1325 1330 1335	Ala Leu Tyr Gln Ala His Leu Ala Met Gly Tyr Ala Pro Ile Ser 1340 1345 1350	Ile Pro Ala Pro Met Arg Asn Asp Asp Leu Val Ser Ile Ala Ala 1355 1360 1365	Thr Tyr Asp Arg Ala Val Ile Ala Tyr Leu Gly Lys Asp Val Pro 1370 1375 1380	Gly Gly Gly Ser Thr Arg Ile Thr Arg His Asp Glu Gly Ala Gly 1385 1390 1395	Ser Val Val Ser Phe Asp Tyr Asn Ala Ala Val Gln Ala Ser Ala 1400 1405 1410	Val Pro Arg Leu Gly Gln Val Tyr Val Leu Ile Ser Asn Asp Gly 1415 1420 1425	Gln Gly Ala Arg Ala Val Leu Leu Ala Glu Asp Leu Ala Trp Ala 1430 1435 1440	Gly Ser Gly Ser Ala Leu Asp Val Leu Asn Glu Arg Leu Val Thr 1445 1450 1455	Leu Phe Pro Ala Pro Val 1460
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---------------------------------

<210> SEQ ID NO 3

<211> LENGTH: 403

<212> TYPE: PRT

-continued

<213> ORGANISM: Bordetella pertussis

<400> SEQUENCE: 3

Met Ala Pro Leu Leu Val Leu Gly Phe Ala Ser Gly Leu Pro Leu Ala
 1 5 10 15
 Leu Ser Ser Gly Thr Leu Gln Ala Trp Ala Thr Val Glu Asn Val Ser
 20 25 30
 Leu Gln Ser Ile Gly Phe Leu Thr Leu Ala Gly Thr Ala Tyr Thr Leu
 35 40 45
 Lys Phe Leu Trp Ala Pro Leu Ile Asp Arg Tyr Val Pro Pro Phe Leu
 50 55 60
 Gly Arg Arg Arg Gly Trp Met Leu Leu Thr Gln Val Leu Leu Ala Ala
 65 70 75 80
 Ala Ile Met Val Met Gly Met Leu Ser Pro Gly Ser Ala Leu Leu Pro
 85 90 95
 Leu Ala Leu Val Ala Val Leu Val Ala Phe Leu Ser Ala Ser Gln Asp
 100 105 110
 Ile Ala Phe Asp Ala Tyr Ser Thr Asp Val Leu Arg Gln Glu Glu Arg
 115 120 125
 Gly Ala Gly Ala Ala Met Arg Val Met Gly Tyr Arg Leu Ala Met Ile
 130 135 140
 Val Ser Gly Gly Leu Ala Leu Ile Val Ala Asp Arg Trp Leu Gly Trp
 145 150 155 160
 Gly Asn Thr Tyr Val Leu Met Gly Gly Leu Met Leu Ala Cys Ala Leu
 165 170 175
 Gly Thr Leu Trp Ala Pro Glu Pro Glu Arg Pro Ala Asn Pro Pro Arg
 180 185 190
 Asp Leu Gly Ala Ala Val Val Glu Pro Phe Arg Glu Phe Phe Ser Arg
 195 200 205
 Arg Gly Ala Ile Asp Met Leu Leu Leu Ile Val Leu Tyr Lys Leu Gly
 210 215 220
 Asp Ala Phe Ala Gly Ala Leu Ser Thr Thr Phe Leu Leu Arg Gly Ala
 225 230 235 240
 Gly Phe Ser Ala Thr Glu Val Gly Thr Val Asn Lys Val Leu Gly Leu
 245 250 255
 Ala Ala Thr Ile Val Gly Ala Leu Ala Gly Gly Ser Ile Met Thr Arg
 260 265 270
 Trp Gly Leu Tyr Arg Ser Leu Met Ala Phe Gly Leu Leu Gln Ala Val
 275 280 285
 Ser Asn Leu Gly Tyr Trp Leu Ile Ala Val Ser Pro Lys Asn Leu Tyr
 290 295 300
 Leu Met Gly Leu Ala Val Gly Val Glu Asn Leu Cys Gly Gly Leu Gly
 305 310 315 320
 Thr Ala Ser Phe Val Ala Leu Leu Met Ala Met Cys Arg Gln Gln Phe
 325 330 335
 Ser Ala Thr Gln Phe Ala Leu Leu Ser Ala Leu Ala Ala Val Gly Arg
 340 345 350
 Thr Tyr Leu Ala Gly Pro Leu Thr Pro Val Leu Val Glu Trp Leu Asp
 355 360 365
 Trp Pro Gly Phe Phe Ile Val Thr Val Leu Ile Ala Leu Pro Gly Leu
 370 375 380
 Trp Leu Leu Arg Leu Arg Arg Asn Val Ile Asp Glu Leu Asp Ala Gln
 385 390 395 400

-continued

Thr Ala Arg

<210> SEQ ID NO 4

<211> LENGTH: 491

<212> TYPE: PRT

<213> ORGANISM: Escherichia coli

<400> SEQUENCE: 4

```

Met Ser Ser Gln Tyr Leu Arg Ile Phe Gln Gln Pro Arg Ser Ala Ile
 1           5           10           15
Leu Leu Ile Leu Gly Phe Ala Ser Gly Leu Pro Leu Ala Leu Thr Ser
 20           25           30
Gly Thr Leu Gln Ala Trp Met Thr Val Glu Asn Ile Asp Leu Lys Thr
 35           40           45
Ile Gly Phe Phe Ser Leu Val Gly Gln Ala Tyr Val Phe Lys Phe Leu
 50           55           60
Trp Ser Pro Leu Met Asp Arg Tyr Thr Pro Pro Phe Phe Gly Arg Arg
 65           70           75           80
Arg Gly Trp Leu Leu Ala Thr Gln Ile Leu Leu Leu Val Ala Ile Ala
 85           90           95
Ala Met Gly Phe Leu Glu Pro Gly Thr Gln Leu Arg Trp Met Ala Ala
 100          105          110
Leu Ala Val Val Ile Ala Phe Cys Ser Ala Ser Gln Asp Ile Val Phe
 115          120          125
Asp Ala Trp Lys Thr Asp Val Leu Pro Ala Glu Glu Arg Gly Ala Gly
 130          135          140
Ala Ala Ile Ser Val Leu Gly Tyr Arg Leu Gly Met Leu Val Ser Gly
 145          150          155          160
Gly Leu Ala Leu Trp Leu Ala Asp Lys Trp Leu Gly Trp Gln Gly Met
 165          170          175
Tyr Trp Leu Met Ala Ala Leu Leu Ile Pro Cys Ile Ile Ala Thr Leu
 180          185          190
Leu Ala Pro Glu Pro Thr Asp Thr Ile Pro Val Pro Lys Thr Leu Glu
 195          200          205
Gln Ala Val Val Ala Pro Leu Arg Asp Phe Phe Gly Arg Asn Asn Ala
 210          215          220
Trp Leu Ile Leu Leu Leu Ile Val Leu Tyr Lys Leu Gly Asp Ala Phe
 225          230          235          240
Ala Met Ser Leu Thr Thr Thr Phe Leu Ile Arg Gly Val Gly Phe Asp
 245          250          255
Ala Gly Glu Val Gly Val Val Asn Lys Thr Leu Gly Leu Leu Ala Thr
 260          265          270
Ile Val Gly Ala Leu Tyr Gly Gly Ile Leu Met Gln Arg Leu Ser Leu
 275          280          285
Phe Arg Ala Leu Leu Ile Phe Gly Ile Leu Gln Gly Ala Ser Asn Ala
 290          295          300
Gly Tyr Trp Leu Leu Ser Ile Thr Asp Lys His Leu Tyr Ser Met Gly
 305          310          315          320
Ala Ala Val Phe Phe Glu Asn Leu Cys Gly Gly Met Gly Thr Ser Ala
 325          330          335
Phe Val Ala Leu Leu Met Thr Leu Cys Asn Lys Ser Phe Ser Ala Thr
 340          345          350
Gln Phe Ala Leu Leu Ser Ala Leu Ser Ala Val Gly Arg Val Tyr Val
 355          360          365

```

-continued

Gly Pro Val Ala Gly Trp Phe Val Glu Ala His Gly Trp Ser Thr Phe
 370 375 380

Tyr Leu Phe Ser Val Ala Ala Val Pro Gly Leu Ile Leu Leu Leu
 385 390 395 400

Val Cys Arg Gln Thr Leu Glu Tyr Thr Arg Val Asn Asp Asn Phe Ile
 405 410 415

Ser Arg Thr Glu Tyr Pro Ala Gly Tyr Ala Phe Ala Met Trp Thr Leu
 420 425 430

Ala Ala Gly Ile Ser Leu Leu Ala Val Trp Leu Leu Leu Leu Thr Met
 435 440 445

Asp Ala Leu Asp Leu Thr His Phe Ser Phe Leu Pro Ala Leu Leu Glu
 450 455 460

Val Gly Val Leu Val Ala Leu Ser Gly Val Val Leu Gly Gly Leu Leu
 465 470 475 480

Asp Tyr Leu Ala Leu Arg Lys Thr His Leu Met
 485 490

<210> SEQ ID NO 5
 <211> LENGTH: 32
 <212> TYPE: DNA
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Oligonucleotide A

<400> SEQUENCE: 5

tataaatcga tattcctgct ggtttcggtc tc

32

<210> SEQ ID NO 6
 <211> LENGTH: 30
 <212> TYPE: DNA
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Oligonucleotide B

<400> SEQUENCE: 6

tatagctagc aagttgggaa acgacaccac

30

<210> SEQ ID NO 7
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Oligonucleotide C

<400> SEQUENCE: 7

taagaagcaa aataagccag gcatt

25

<210> SEQ ID NO 8
 <211> LENGTH: 30
 <212> TYPE: DNA
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Oligonucleotide D

<400> SEQUENCE: 8

tataccatgg cgccgctgct ggtgctgggc

30

<210> SEQ ID NO 9
 <211> LENGTH: 30
 <212> TYPE: DNA
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Oligonucleotide E

-continued

<400> SEQUENCE: 9

tatatctaga cgctggccgt aaccttagca 30

<210> SEQ ID NO 10

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Oligonucleotide F

<400> SEQUENCE: 10

tatagaattc gctcggttcg ctggtcaagg 30

<210> SEQ ID NO 11

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Oligonucleotide G

<400> SEQUENCE: 11

tatatctaga gcaatgccga ttcatttcta 30

<210> SEQ ID NO 12

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Oligonucleotide H

<400> SEQUENCE: 12

tatatctaga gcggccttta ttgcttttcc 30

<210> SEQ ID NO 13

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Oligonucleotide I

<400> SEQUENCE: 13

tataaagctt ctcatgcacg ccggcttctc 30

What is claimed is:

1. A method of making a live attenuated *Bordetella* strain for use in a vaccine, the method comprising the steps of:

providing a live parent *Bordetella* strain comprising a pertussis toxin gene (ptx), a dermonecrotic toxin gene (dnt) and a *Bordetella* ampG gene;

mutating the ptx in the live parent *Bordetella* strain;

deleting or mutating the dnt gene in the live parent *Bordetella* strain; and

replacing the *Bordetella* ampG gene in the live parent *Bordetella* strain with a heterologous ampG gene, wherein the resulting live mutated *Bordetella* strain is able to colonize the respiratory tract of a subject and induce protective immunity in the subject.

2. The method of claim 1, wherein the step of replacing the *Bordetella* ampG gene in the live parent *Bordetella* strain with the heterologous ampG gene causes the live attenuated *Bordetella* strain to express less than 5% of the tracheal cytotoxin (TCT) activity compared to the TCT activity expressed by the live parent *Bordetella* strain.

3. The method of claim 1, further comprising the step of introducing into the live attenuated *Bordetella* strain a heterologous expression platform which carries a heterologous antigen.

4. The method of claim 3, wherein the heterologous antigen is one expressed by a respiratory pathogen selected from the group consisting of: *Neisseria*, *Pneumophila*, *Yersinia*, *Pseudomonas*, *Mycobacteria*, and *Influenza*.

* * * * *